

Clinical features and virological confirmation of perinatal dengue infection in Jambi, Indonesia- A case report

by Herlambang Herlambang

Submission date: 21-Apr-2021 11:23AM (UTC+0700)

Submission ID: 1565346686

File name: erinatal_dengue_infection_in_Jambi,_Indonesia-_A_case_report.pdf (834.74K)

Word count: 3043

Character count: 16557



Contents lists available at ScienceDirect

International Journal of Infectious Diseases

journal homepage: www.elsevier.com/locate/ijid



Case Report

Clinical features and virological confirmation of perinatal dengue infection in Jambi, Indonesia: A case report



Sotianingsih Haryanto^a, Benediktus Yohan^b, Marsha Sinditia Santoso^b, Rahma F. Hayati^b,
Dionisius Denis^b, Gorga I.V.W. Udjung^a, Herlambang^c, R. Indah Kendarsari^b,
Hidayat Trimarsanto^b, R. Tedjo Sasmono^{b,*}

^a *10 m Hospital and Raden Matta Her Hospital, Jambi, Indonesia*

^b *Eijkman Institute for Molecular Biology, Ministry of Research, Technology, and Higher Education of the Republic of Indonesia, Jakarta, Indonesia*

^c *Faculty of Medicine and Health Science, Universitas Jambi, Indonesia*

ARTICLE INFO

Article history:

Received 15 May 2019

Received in revised form 18 July 2019

Accepted 22 July 2019

Corresponding Editor: Eskild Petersen, Aarhus, Denmark

Keywords:

Dengue

Pregnancy

Perinatal

Vertical transmission

Neonate

Next-generation sequencing

ABSTRACT

The hyper-endemicity of dengue in Indonesia poses a significant threat of dengue virus (DENV) vertical transmission during pregnancy. A 29-year-old female at 38 weeks of pregnancy presented to hospital with acute fever and later confirmed with DENV infection. Due to signs of fetal distress, the neonate was delivered by emergency caesarean section. The mother developed a dengue critical phase post-caesarean with excessive bleeding and required blood transfusion. During the 6th day of life, the neonate was diagnosed and later confirmed with dengue. Next-generation sequencing of DENV RNA isolated directly from sera of both mother and neonate revealed identical DENV-2 whole-genome sequences. Plaque reduction neutralization test (PRNT) detected anti-dengue antibodies in both mother and neonate. Altogether, our data confirmed the occurrence of vertical transmission. Dengue vertical transmission during pregnancy may lead to severe manifestation, hence early diagnosis, close monitoring, and prompt intervention are critical.

© 2019 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Background

13 Dengue is a growing global health problem with an estimated 390 million infections per year worldwide (Bhatt et al., 2013). Dengue has been found in increasing numbers in all 34 provinces in Indonesia, posing significant risk to 250 million population and placing a high economic and disease burden on the country (Wahyono et al., 2017).

Dengue virus (DENV) infection can occur in any age group, including pregnant women and their neonates. The DENV-infected pregnant women are considered to be a vulnerable group, having a higher risk of developing severe dengue, hemorrhagic complications, and death, though the mechanisms are still unclear (Charlier et al., 2017). Here, we report a case of symptomatic maternal dengue infection and subsequent symptomatic disease in the

neonate confirmed through whole-genome sequencing with evidence of anti-DENV antibody transfer, suggesting vertical transmission.

14 Case presentation

A 29-year-old female was admitted to Siloam Hospital, Jambi, Indonesia with symptoms of fever that began 3 days prior, joint and muscle pains, headache, and loss of appetite. She was 38 weeks pregnant with her second child. She had a spontaneous vaginal delivery for her first pregnancy. Her vitals were normal upon admission and she tested positive for DENV NS1 antigen with a platelet count of 149,000/ μ L. Serological testing using Panbio Dengue Duo IgG/IgM rapid tests resulted positive for anti-DENV IgG and IgM and further confirmed using Panbio Dengue Duo IgG/IgM ELISA. Other laboratory results were normal. The fetus showed a fetal heart rate of 170 bpm, demonstrating fetal distress. Due to continued signs of fetal distress observed as increased fetal heart rate and reduced movement, the patient underwent emergency caesarean section the next day.

A female neonate weighing 2.9 kg was delivered. The amniotic fluid was stained with meconium and the neonate displayed signs

Abbreviations: BWA, Burrows-Wheeler Alignment; BPM, beats per minute; DENV, dengue virus; ICU, intensive care unit; NGS, next-generation sequencing; NS1, non-structural protein 1.

* Corresponding author at: Eijkman Institute for Molecular Biology, Jl. Diponegoro 69, Jakarta 10430, Indonesia.

E-mail address: sasmono@eijkman.go.id (R. T. Sasmono).

18
<https://doi.org/10.1016/j.ijid.2019.07.019>

1201-9712/© 2019 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

of respiratory distress at birth with 1st- and 5th-minute Apgar scores of 6 and 8, respectively. The neonate's vital signs returned to normal and had normal lab results during the next 4 days of life. During the caesarean section, the mother lost 200 mL of blood. Post-caesarean section, her platelets began decreasing, reaching their lowest on the 2nd day post-caesarian section at 59,000/ μ L. On the 6th day post-caesarian, her symptoms and laboratory results returned to normal and she was discharged from the hospital two days later.

On the neonate's 6th day of life, her temperature rose to 38.6 °C and she began showing signs of respiratory distress with increased respiratory rate and cyanotic extremities. She was then admitted to neonatal ICU. There were no signs of mosquito bite in this specialized containment. Laboratory results ¹⁷ demonstrated a plummeting platelet count to 23,000/ μ L and on the 2nd day of

fever, she tested positive for DENV NS1 antigen. The platelet count continued to drop as far as 11,000/ μ L and she received a total of 150 mL of thrombocyte concentrate transfusion spread over the next 3 days. Serological test using Panbio anti-DENV IgG/IgM rapid assay results were negative. However, when we tested using ELISA, the results were positive only for anti-DENV IgM. There were no signs of hemorrhage, pleural effusion, or ascites. On the 10th day of life and 5th day post-fever onset, the platelet count returned to normal at 256,000/ μ L. She was discharged from the hospital the next day.

During the observation, blood samples were collected from both mother and neonate during their viremic phase at the third day of fever for the mother and second day of fever for the ¹²neate. Serum samples were separated from venous blood and used for RNA extraction using QIAamp Viral RNA Mini kit (Qiagen). An



Figure 1. Maximum Clade Credibility (MCC) tree of DENV-2 Cosmopolitan ¹genotype grouping generated by Bayesian inference method as implemented in BEAST using TN93 evolution model, gamma parameter rates, and molecular clock analysis from the Envelope gene sequences. The Jambi perinatal isolates (red font) were grouped into a Jambi-Sumatran clade (highlighted ¹⁶ together with other isolates from Sumatra 2009–2010 and Jambi 2015 (blue font). The posterior probabilities of the clades were indicated as numbers in the node labels. Whole-genome sequences of the DENV-2 isolates have been deposited in GenBank with accession number MK411558 and MK411559.

initial DENV detection and serotyping was performed using Simplexa Dengue real-time RT-PCR (Sasmono et al., 2014) and revealed the presence of dengue virus serotype 2 (DENV-2) in both samples. An in-depth analysis was performed using next-generation sequencing (NGS) where RNAs were extracted from 6×10^4 μ L of serum sample, pooled, and used as template in NGS library preparation using TruSeq RNA library preparation kit (Illumina) using a metagenomic approach. Library was run in Illumina MiSeq NGS system and the resulting reads processed using BWA alignment and SPAdes de novo assembly bioinformatic softwares. The DENV-2 was detected in 0.14% and 0.02% of total sequence reads with $75\times$ and $20\times$ depth for mother and neonate samples, respectively. A total of 95% DENV genome coverage was achieved for both samples. Sequence identity analysis of the genome revealed a 100% identity (GenBank accession numbers MK411558 and MK411559). The DENV sequences were genotyped as Cosmopolitan, according to Twiddy, et al. (Twiddy et al., 2002) (Figure 1).

The titer of anti-DENV antibody in serum samples was measured using a standard dengue Plaque Reduction Neutralization Test (PRNT) (Timiryasova et al., 2013) where 2-fold serially diluted serum samples was performed and challenged with the parental strains of the recombinant CYD vaccine viruses from all four DENV serotypes (Sasmono et al., 2018). Serum samples from the mother showed a detectable level of antibodies to all four serotypes in high levels. A similar profile, at a much lesser degree, was observed in the neonate's serum (Supplementary Table).

Discussion

In this case report, the mother was admitted to hospital with mild dengue symptoms which later developed into severe manifestation. The confirmation of DENV infection using NS1 antigen detection increased the awareness of the medical staff to monitor the fetal distress status which led to the decision to conduct caesarean section delivery. The elective decision to safeguard the fetus from potential harm of dengue-related risks has been described previously (Chye et al., 1997), although hazards of surgical intervention in patients with acute DENV infection are possible (Thaithumyanon et al., 1994).

Severe manifestations of dengue in the mother were observed within 3 days post-caesarean section, with evident thrombocytopenia and blood loss requiring blood transfusion, as has been observed elsewhere (Chye et al., 1997). The neonate showed evidence of dengue infection in her 6th day, with thrombocytopenia and positive DENV-NS1 antigen detection. However, with careful monitoring and intervention, the mother and neonate's conditions were improved and did not last to long-term sequelae. The clinical manifestations of the DENV-infected neonates are variable, from asymptomatic disease, typical dengue symptoms such as fever and rash, signs of plasma leakage such as ascites and pleural effusion, to dengue shock syndrome (Sinhobahu et al., 2014; Yang et al., 2015; Arragain et al., 2017). Dengue symptoms in the mother, obstetric complications associated with dengue, and clinical outcomes of the neonate have been described in a systematic review on maternal dengue and pregnancy outcomes (Pouliot et al., 2010).

The occurrence of dengue mother-to-child transmission has been confirmed through the detection of IgM/IgG antibodies, NS1 antigen, viral isolation, and nucleic acid (Sinhobahu et al., 2014; Arragain et al., 2017; Ribeiro et al., 2013). We performed a metagenomic NGS approach to sequence the whole genome of the DENV. The genomes sequences derived from both mother and neonate showed identical sequences, inferring the vertical transmission of DENV. Phylogenetic analysis revealed that the DENV-2 was classified into Cosmopolitan genotype and

closely-related to strains circulated in Jambi (Haryanto et al., 2016). To our knowledge, our study is the first to confirm dengue perinatal transmission using NGS technology.

The antibody profiles of both mother and neonate were highly similar showing evidence of antibody transfer from mother to neonate. Low titer of antibodies (DENV neutralizing antibodies) in the neonate may be the cause of negative IgG testing using rapid test and ELISA, which may be explained by the lower sensitivity of the assays compared to PRNT. It was also of note that the mother was likely to acquire secondary infection, shown by the presence of antibodies to all four DENV serotypes. The presence of all four anti-DENV antibodies in the neonate confirmed the occurrence of antibody transfer, since natural DENV infection in the neonate would be likely to generate primary infection and antibody to the particular serotype. Studies reported greater levels of anti-DENV antibodies in infants due to vertical transmission (Chau et al., 2009; Castanha et al., 2016; Kliks et al., 1988). However, the lower level of IgG transferred to the neonate may be explained by the findings that the transfer of total IgG is decreased as the level of maternal IgG increased (Castanha et al., 2016) as observed by the high levels of maternal IgG in our study (Supplementary Table).

Although cord blood analysis was not performed, the evidence of positive DENV NS1 antigen in both mother and neonate and the identical DENV-2 genome sequences derived from both specimens indicated the occurrence of vertical transmission. The evidence of mosquito bite was ruled out since the mother and neonate were cared for in the ICU with containment from pathogens and germs, including mosquito. The onset of fever within 7 days of onset also supported the diagnosis of vertical transmission, as also observed in other case reports (Phupong, 2001; Bopeththa et al., 2018).

In conclusion, we report the dengue vertical transmission and confirmed it using DENV NS1 antigen, and DENV RNA detection (and whole-genome sequencing), and anti-DENV IgG/IgM antibodies. Dengue poses significant risks to women in pregnancy and the apparent vertical transmission to neonates should be taken into account while incorporating a vigilant disease management.

Authors' contributions

H, RTS set the conceptual design of the study; SH, GIVWU, MSS, H performed clinical care, diagnosis, data collection, and analysis on patients' status. BY, RFH, DD, RIK performed the experiments and data analysis; HT analyzed the NGS data and performed bioinformatic analyses; RTS, SH, BY, MSS made the first draft of the manuscript; RTS contributed to the generation of final manuscript.

Consent for publication

Written informed consent was obtained from the patient for publication of this case report.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We would like to express gratitude to all staff of Siloam Hospital, Jambi involved in patients' care and management and to Pandu Biosains for support in providing Illumina NGS reagents. This study was funded by APBN grant from the Ministry of Research, Technology, and Higher Education of the Republic Indonesia to Dengue Research Unit of the Eijkman Institute.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.ijid.2019.07.019>.

References

- Arragain L, Dupont-Rouzeyrol M, O'Connor O, Sigur N, Grangeon J-P, Huguon E, et al. Vertical transmission of dengue virus in the peripartum period and viral kinetics in newborns and breast milk: new data. *J Pediatr Infect Dis Soc* 2017;6:324–31, doi:<http://dx.doi.org/10.1093/jpids/piw058>.
- Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and burden of dengue. *Nature* 2013;496:504–7, doi:<http://dx.doi.org/10.1038/nature12060>.
- Bopeththa BVKM, Hemapriya S, Gayan Niranga KK, Kotigala DSK. A case report of dengue haemorrhagic fever during the peripartum period: challenges in management and a case of vertical dengue transmission. *BMC Infect Dis* 2018;18:427, doi:<http://dx.doi.org/10.1186/s12879-018-3352-x>.
- Castanha PMS, Braga C, Cordeiro MT, Souza AI, Silva CD, Martelli CMT, et al. Placental transfer of dengue virus (DENV)-specific antibodies and kinetics of DENV infection-enhancing activity in Brazilian infants. *J Infect Dis* 2016;214:265–72, doi:<http://dx.doi.org/10.1093/infdis/jiw143>.
- Charlier C, Beaudoin M-C, Coudert T, Lortholary O, Lecuit M. Arboviruses and pregnancy: maternal, fetal, and neonatal effects. *Lancet Child Adolesc Health* 2017;1:134–46, doi:[http://dx.doi.org/10.1016/S2352-4642\(17\)30021-4](http://dx.doi.org/10.1016/S2352-4642(17)30021-4).
- Chau TNB, Hieu NT, Anders KL, Wolbers M, Le Bich L, Lu Thi Minh H, et al. Dengue virus infections and maternal antibody decay in a prospective birth cohort study of Vietnamese Infants. *J Infect Dis* 2009;200:1893–900, doi:<http://dx.doi.org/10.1086/648407>.
- Chye JK, Lim CT, Ng KB, Lim JM, George R, Lam SK. Vertical transmission of dengue. *Clin Infect Dis* 1997;25:1374–7.
- Haryanto S, Hayati RF, Yohan B, Sjabat L, Sihite IF, Fahri S, et al. The molecular and clinical features of dengue during outbreak in Jambi, Indonesia in 2015. *Pathog Glob Health* 2016;110:119–29, doi:<http://dx.doi.org/10.1080/20477724.2016.1184864>.
- Kliks SC, Nimmanitya S, Nisalak A, Burke DS. Evidence that maternal dengue antibodies are important in the development of dengue hemorrhagic fever in infants. *Am J Trop Med Hyg* 1988;38:411–9.
- Phupong V. Dengue fever in pregnancy: a case report. *BMC Pregnancy Childbirth* 2001;1:7.
- Pouliot SH, Xiong X, Harville E, Paz-Soldan V, Tomashek KM, Breart G, et al. Maternal dengue and pregnancy outcomes: a systematic review. *Obstet Gynecol Surv* 2010;65:107–18, doi:<http://dx.doi.org/10.1097/OGX.0b013e3181cb8fbc>.
- Ribeiro CF, Lopes VGS, Brasil P, Coelho J, Muniz AG, Nogueira RMR. Perinatal transmission of dengue: a report of 7 cases. *J Pediatr* 2013;163:1514–6, doi:<http://dx.doi.org/10.1016/j.jpeds.2013.06.040>.
- Sasmono RT, Aryati A, Wardhani P, Yohan B, Trimarsanto H, Fahri S, et al. Performance of Simplex dengue molecular assay compared to conventional and SYBR green RT-PCR for detection of dengue infection in Indonesia. *PLoS One* 2014;9:e103815, doi:<http://dx.doi.org/10.1371/journal.pone.0103815>.
- Sasmono RT, Taurel A-F, Prayitno A, Sitompul H, Yohan B, Hayati RF, et al. Dengue virus serotype distribution based on serological evidence in pediatric urban population in Indonesia. *PLoS Negl Trop Dis* 2018;12:e0006616, doi:<http://dx.doi.org/10.1371/journal.pntd.0006616>.
- Sinhahabu VP, Sathananthan R, Malavige GN. Perinatal transmission of dengue: a case report. *BMC Res Notes* 2014;7:, doi:<http://dx.doi.org/10.1186/1756-0500-7-795>.
- Thaithumyanon P, Thisyakorn U, Deerojanawong J, Innis BL. Dengue infection complicated by severe hemorrhage and vertical transmission in a parturient woman. *Clin Infect Dis* 1994;18:248–9.
- Timiryasova TM, Bonaparte MI, Luo P, Zedar R, Hu BT, Hildreth SW. Optimization and validation of a plaque reduction neutralization test for the detection of neutralizing antibodies to four serotypes of dengue virus used in support of dengue vaccine development. *Am J Trop Med Hyg* 2013;88:962–70, doi:<http://dx.doi.org/10.4269/ajtmh.12-0461>.
- Twiddy SS, Farrar JJ, Vinh Chau N, Wills B, Gould EA, Gritsun T, et al. Phylogenetic relationships and differential selection pressures among genotypes of dengue-2 virus. *Virology* 2002;298:63–72.
- Wahyono TYM, Nealon J, Beucher S, Prayitno A, Moureau A, Nawawi S, et al. Indonesian dengue burden estimates: review of evidence by an expert panel. *Epidemiol Infect* 2017;145:2324–9, doi:<http://dx.doi.org/10.1017/S0950268817001030>.
- Yang S-T, Chen H-L, Yeh C-T, Lee W-T. Vertical transmission of dengue fever: first case reported in Taiwan. *J Formos Med Assoc Taiwan Yi Zhi* 2015;114:558–9, doi:<http://dx.doi.org/10.1016/j.jfma.2015.02.005>.

Clinical features and virological confirmation of perinatal dengue infection in Jambi, Indonesia- A case report

ORIGINALITY REPORT

19%
SIMILARITY INDEX

13%
INTERNET SOURCES

15%
PUBLICATIONS

6%
STUDENT PAPERS

PRIMARY SOURCES

1	Puspa Wardhani, Aryati Aryati, Benediktus Yohan, Hidayat Trimarsanto, Tri Y. Setianingsih, Dwiyantri Puspitasari, Muhammad Vitanata Arfijanto, Bramantono Bramantono, Suharto Suharto, R. Tedjo Sasmono. "Clinical and virological characteristics of dengue in Surabaya, Indonesia", PLOS ONE, 2017 Publication	3%
2	Submitted to The University of Memphis Student Paper	2%
3	www.scribd.com Internet Source	2%
4	bmcpulmmed.biomedcentral.com Internet Source	1%
5	curis.ku.dk Internet Source	1%
6	B. V. K. M. Bopeththa, Sardha Hemapriya, K. K. Gayan Niranga, Danushka S. K. Kotigala. "A	1%

case report of dengue haemorrhagic fever during the peripartum period: challenges in management and a case of vertical dengue transmission", BMC Infectious Diseases, 2018

Publication

-
- | | | |
|---|--|---|
| <div style="background-color: #8B4513; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">7</div> | <div style="color: #8B4513; font-weight: bold; font-size: 1.2em;">f1000research.com</div> <div style="color: #8B4513; font-size: 0.9em;">Internet Source</div> | <div style="font-size: 2.5em; font-weight: bold;">1</div> % |
|---|--|---|
-
- | | | |
|---|--|---|
| <div style="background-color: #0056b3; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">8</div> | <div style="color: #0056b3; font-weight: bold; font-size: 1.2em;">www.asianpids.org</div> <div style="color: #0056b3; font-size: 0.9em;">Internet Source</div> | <div style="font-size: 2.5em; font-weight: bold;">1</div> % |
|---|--|---|
-
- | | | |
|---|---|---|
| <div style="background-color: #800080; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">9</div> | <div style="color: #800080; font-weight: bold; font-size: 1.2em;">Breno Lima de Almeida, Daniel Vicente Silva, Mateus Santana do Rosário, Bruno Simões Dias Gonçalves et al. "Biliary Ascariasis and severe bacterial outcomes: Report of three cases from a Pediatric Hospital in Brazil", International Journal of Infectious Diseases, 2020</div> <div style="color: #800080; font-size: 0.9em;">Publication</div> | <div style="font-size: 2.5em; font-weight: bold;">1</div> % |
|---|---|---|
-
- | | | |
|--|---|---|
| <div style="background-color: #6b8e23; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">10</div> | <div style="color: #6b8e23; font-weight: bold; font-size: 1.2em;">innovareacademics.in</div> <div style="color: #6b8e23; font-size: 0.9em;">Internet Source</div> | <div style="font-size: 2.5em; font-weight: bold;">1</div> % |
|--|---|---|
-
- | | | |
|--|--|---|
| <div style="background-color: #000080; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">11</div> | <div style="color: #000080; font-weight: bold; font-size: 1.2em;">Keita Suzuki, Juthamas Phadungsombat, Emi E. Nakayama, Akatsuki Saito et al. "Genotype replacement of dengue virus type 3 and clade replacement of dengue virus type 2 genotype Cosmopolitan in Dhaka, Bangladesh in 2017", Infection, Genetics and Evolution, 2019</div> <div style="color: #000080; font-size: 0.9em;">Publication</div> | <div style="font-size: 2.5em; font-weight: bold;">1</div> % |
|--|--|---|
-
- | | | |
|--|---|--|
| <div style="background-color: #0070c0; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">12</div> | <div style="color: #0070c0; font-weight: bold; font-size: 1.2em;">www.ncbi.nlm.nih.gov</div> <div style="color: #0070c0; font-size: 0.9em;">Internet Source</div> | |
|--|---|--|

<1 %

13

Fu, Yilong, Yen-Liang Chen, Maxime Herve, Feng Gu, Pei-Yong Shi, and Francesca Blasco. "Development of a FACS-based assay for evaluating antiviral potency of compound in dengue infected peripheral blood mononuclear cells", Journal of Virological Methods, 2013.

Publication

<1 %

14

balimedicaljournal.org

Internet Source

<1 %

15

Advances in Microbial Ecology, 2012.

Publication

<1 %

16

Akhtar Ali, Ijaz Ali. "The Complete Genome Phylogeny of Geographically Distinct Dengue Virus Serotype 2 Isolates (1944-2013) Supports Further Groupings within the Cosmopolitan Genotype", PLOS ONE, 2015

Publication

<1 %

17

Sri Masyeni, Marsha S. Santoso, Putu Dyah Widyaningsih, DG Wedha Asmara, Firzan Nainu, Harapan Harapan, R Tedjo Sasmono. "Serological cross-reaction and coinfection of dengue and COVID-19 in Asia: Experience from Indonesia", International Journal of Infectious Diseases, 2021

Publication

<1 %

18	dspace.alquds.edu Internet Source	<1 %
19	www.biorxiv.org Internet Source	<1 %
20	Endang S. Kusmintarsih, Rahma F. Hayati, Oktaviani N. Turnip, Benediktus Yohan et al. "Molecular characterization of dengue viruses isolated from patients in Central Java, Indonesia", Journal of Infection and Public Health, 2017 Publication	<1 %
21	repositorio.ufpe.br Internet Source	<1 %
22	soroylardo.com Internet Source	<1 %
23	www.nature.com Internet Source	<1 %
24	www.pure.ed.ac.uk Internet Source	<1 %
25	Minh Huong Phu Ly, Meng Ling Moi, Thi Bich Hau Vu, Mya Myat Ngwe Tun et al. "Dengue virus infection-enhancement activity in neutralizing antibodies of healthy adults before dengue season as determined by using FcyR-expressing cells", BMC Infectious Diseases, 2018	<1 %

26

Ying Wang, Lulu Si, Yayan Luo, Xiaolan Guo, Junmei Zhou, Danyun Fang, Huijun Yan, Gucheng Zeng, Lifang Jiang. "Replacement of pr gene with Japanese encephalitis virus pr using reverse genetics reduces antibody-dependent enhancement of dengue virus 2 infection", Applied Microbiology and Biotechnology, 2015

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Clinical features and virological confirmation of perinatal dengue infection in Jambi, Indonesia- A case report

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4