

Artikel

by Madyawati Latief

Submission date: 29-Mar-2021 11:35AM (UTC+0700)

Submission ID: 1545092854

File name: th_Cap-e_Methods_to_Overcome_Oxidative_Stress_in_Thalasemia.pdf (813.39K)

Word count: 2133

Character count: 12047

Antioxidant Activity from Perepat Plant (*Sonneratia alba*) Ethanol Leaf Extract with Cap-e Methods to Overcome Oxidative Stress in Thalassemia

Madyawati Latief¹, Andita Utami¹, Nurul Fadhillah¹, Restina Bemis², Hilda Amanda², Heriyanti², Martina Asti Rahayu², Yusnaidar^{1,2}, Wilda Syahr², Muhaimin^{1,2,*}

4

¹Department of Chemistry, Faculty of Science and Technology, University of Jambi, Jambi, 36361, Indonesia

²Department of Chemistry Education, Faculty of Education, University of Jambi, Jambi, 36361 Indonesia

Abstract

Thalassemia is a genetic disease with symptoms of anemia as a result of partial or complete disorder of the hemoglobin synthesis in red blood cells. Consumption of blood supplementation and routine blood transfusion to overcome the anemia symptoms in thalassemia patients can lead to excessive absorption of iron (Fe) content that can trigger the occurrence of oxidative stress. Therefore, antioxidant supplements are also important for thalassemia patients. Natural antioxidants derived from plants and fruits can be used to meet the needs of antioxidants in thalassemia patients. The diversity of mangrove plants has the potential to be explored and tested for its bioactivity, one of which is the herbaceous plant (*Sonneratia alba*), which has the potential as a source of antioxidant compounds. Preliminary test of antioxidant activity of ethanol extract has been done by using DPPH method. Furthermore, in-vitro testing with CAP-e (Cellular Antioxidant Protection in erythrocytes) method in normal blood, carrier of thalassemia and thalassemia patient. The results of the test using the ethanol extract from the leaf showed optimum antioxidant activity at a concentration of 300 ppm in normal blood and carrier properties, and 100 ppm in the blood of patients with thalassemia.

Keywords: Mangrove plants, antioxidant, Cap-e, thalassemia

INTRODUCTION

Genetic inherited hemoglobin disorders were originally found in tropical and subtropics [3], but are now commonly found worldwide for migration. The World Health Organization (WHO) has identified hemoglobin disorders as a disease with high impact on public health systems in Western and developing countries. About 7% of the global population is the carrier of the disorder. Among the disorders of hemoglobin, thalassemia clearly contributes (Wheatherall, 2001; Modell and Darlison, 2008; de Franceschi et al., 2011). Due to spontaneous mutations, hemoglobin gene variants are present at low prevalence in all large populations. They fall into two major groups - a structural variant that alters the amino acid sequence resulting in unusual hemoglobin, and thalassemia that decreases or eliminates the production of globin chains (Livingstone, 1985; Modell et al., 2007).

Thalassemia is a genetic disease passed down by Mendel's law in an autosomal recessive manner from parents to his children. This disease is caused by a deficiency of the α and β globin chains that make up hemoglobin (Thein, 2013). Thalassemia is divided into thalassemia minor, intermedia and major. Major thalassemia is the worst type of thalassemia because it can cause severe anemia with ineffective hemolysis and erythropoiesis. Patients suffering from thalassemia major should undergo a monthly blood transfusion to maintain Hb about 9-12 g / dL and increase growth, reduce hepato-splenomegaly and bone deformation (Borgna-Pignati and Galanello, 2004). However, persistent transfusions can cause side effects caused by the accumulation of iron in the organs and cell damage. The deposited iron is responsible for the formation of

reactive oxygen species (ROS) such as superoxide anion (O_2^-),

hydroxyl radical (OH^\cdot), singlet oxygen and hydrogen peroxide (H_2O_2), which induces oxidative stress in thalassemia major patients (Mahdi, 2014). To handle the accumulation of iron, iron sequestering agent is required that serves as an iron binder, but the use of deferoxamine in the long term can cause side effects such as impaired vision, hearing, cardiovascular, digestive, hematology, liver, and nerves (Milena et al., 2010). Because of side effects in the use of iron chelating compounds in the treatment of thalassemia patients, it is necessary alternative therapy that reduces side effects by utilizing natural ingredients that have

antioxidant activity [21]. Antioxidants are protective agents that can quell ROS activity and play an important role in the protection of cells from oxidative damage. Some antioxidant agents such as enzymes (glutathione peroxidase, superoxide dismutase, and catalase), large molecules (ferritin, albumin), small molecules (uric acid, glutathione, bilirubin, ascorbic acid, α -tocopherol, and vitamin E) and some important minerals such as zinc, copper and selenium. Enzymatic antioxidant activity, ie, catalase, glutathione peroxidase and glutathione reductase are found to be drastically reduced in untreated β -thalassemic patients when compared with normal individuals. However, superoxide dismutase activity was found to increase in untreated thalassemia patients when compared with normal individuals (Baumgartner and Van Way, 1999; Dhawan et al., 2005; Simsek et al., 2005).

Based on the literature review, found that vitamin C is reported to inhibit the peroxidation of phospholipid membrane due to oxidative stress in thalassemia patients (Rajagukguk et al., 2014). This suggests that antioxidants are indispensable for treating thalassemia patients. Natural antioxidants derived from plants and fruits can be used to meet the needs of antioxidants in thalassemia patients. Many types of laboratory methods involve red blood cells to evaluate oxidative damage as a result of inflammation, and to evaluate antioxidant protection by natural products (Luqman and Rizvi, 2006). Diversity of plants in Indonesia, especially the diversity of mangrove plants has the potential to be explored its phytochemical content and tested its bioactivity (Herawaty et al., 2011).

One of the plants collected from the mangrove area of Jambi Province, which is a prepat (*Sonneratia alba*) has also been tested for its antioxidant activity against DPPH, shows that ethanol extract provides the best inhibition compared to hexane and acetone extract (Latief et al., 2015). Therefore, in this study was tested in vitro using CAP-e method to see the antioxidant activity of ethanol extract in normal red blood cells, red blood cells of thalassemia carrier and red blood cells of thalassemia patients.

MATERIALS AND METHODS

General: Tools used include: glassware commonly used in chemical laboratories, analytical scales, and rotary evaporator. Meanwhile the tools for CAP-e are vortex, centrifuge, Thermo syringe DVR-3422, Nalgene 0.22 μ M Syringe filter, pipette

(multichannel pipette) and microtips, 96F Nunc Delta black Microwell and Tecan Spectra Fluorescence plate reader.

The solvent used for the extraction of the precipitate leaves is distilled ethanol. In vitro antioxidant testing materials (Cap-e method) are normal red blood cells, red blood cells carrying thalassemia (trait), and thalassemia α and β , saline phosphate buffer (PBS 1X), fetal bovine serum (FBS), ascorbic acid (pse Emsure Merck), ddH₂O, a set of ¹³DA Abuse Cellular ROS Detection Assay kit consisting of 20 mM DCFDA solution (in DMSO), 10X buffer and 55 mM tert-butyl hydrogen peroxide (TBHP) solution.

Plant Materials. Samples of leaf from Perepat plant (*Sonneratia alba*) (Harborne, 1987) as much as 10 kg were collected from mangrove forest area of Kab. Tanjung Jabung Timur, Jambi Province. The leaf sample is cleaned, cut into small pieces, and dried to a constant weight and finely ground.

Extraction and partition. A total of 10 kg of powdered dried leaves of Perepat are macerated with n-hexane solvent, followed by fractionation using acetone solvent and ethanol, to obtain the ethanol fraction of Perepat leaf (*S. alba*) (Muhaimin *et al.*, 2016 and 2017).

Antioxidant activity test by Cap-e method. Test the antioxidant activity of the precursor extract using CAP-e method for normal red blood cells, blood carrier thalassemia and blood thalassemia

patients. Total of 3 mL of fresh blood was centrifuged three times ¹⁸ rate of 2400 rpm in a row for 2.5; 2.5; 10 minutes to separate red blood cells with white blood cells and blood plasma. Furthermore, red blood cells were given treatment using the Abcam DCFDA Cellular ROS Detection Assay (Kang *et al.*, 2010). The order of addition and the number of reagents is adjusted according to the kit instructions. There are 7 variables in test, blanks, negative control, positive control, vitamin C, and 3 variations of concentration from ethanol extract (50 ppm, 100 ppm, and 300 ppm). Blood that has been given each treatment is transferred in dark microplate and stored in dark space at room temperature. After 1 hour then ready to measure the intensity of fluorescence with instrument of Tecan Spectra Fluorescence Plate Reader.

RESULTS AND DISCUSSION

In this study, the leaf part of mangrove plants namely perepat (*S. Alba*) as much as 10 kg has been fractionated. Selected ethanol fractionate for CAP-e test was based on previous research which showed that ethanol extract gave the best value of antioxidant activity compared to hexane and acetone extract (Latief *et al.*, 2015). CAP-e test in normal red blood cells, in this test used 4 samples of normal red blood cells that had previously been screened. Antioxidant activity of ethanol fraction of the leaf shows an optimum free radical inhibition in fractionation of ethanol with 300 ppm constipation.




Figure 1. Graph of antioxidant activity in normal red blood cells.




Figure 2. Graph of Antioxidants Activity in red blood cells from carriers thalassemia.




Figure 3. Graph of antioxidant activity in red blood cells of thalassemia patients.

In ethanol fractionate concentration of 300 ppm, able to reduce the activity of radical bebas, indicated by the low fluorescence value in the measurement. Low fluorescence values indicate at least free radicals that can bind to fluorescence compounds (DCFDA), because free radical activity has been suppressed by antioxidant compounds contained in the ethanol fraction of the leaf.

The CAP-e test on red blood cells of carrier thalassemia. Similarly, the antioxidant activity test in normal red blood cells, antioxidant activity test on red blood cell carrier of thalassemia character also showed the optimum result at fractionation of ethanol with concentration of 300 ppm.

In the carrier of thalassemia trait, the number of red blood cells is less than normal red blood cells. However, its free radical inhibition activity gives a relatively similar pattern.

CAP-e test on red blood cells of thalassemia patients. Antioxidant activity of ethanol fraction in red blood cell of thalassemia patient showed the optimum free radical inhibition at concentration 100 ppm.

In the graph, the ethanol fraction from perepat leaf with a concentration of 300 ppm indicates an increase in fluorescence value. This indicates that at high concentrations, ethanol fractionates act as prooksidan. Thus, free radical inhibition activity that binds to fluorescent compounds is also inhibited. Further research is needed on the optimum concentration of ethanol fraction addition in its activity as an antioxidant in red blood cells of thalassemia patients.

CONCLUSIONS

The results of antioxidant activity test using Cap-e method showed that ethanol extract from mangrove plant from Jambi Province was good, has potency as antioxidant to inhibit oxidative stress found in normal red blood cell, carrier of thalassemia and thalassemia patients. Need further study to know the active compounds that act as antioxidants in the Cap-e method.

5

ACKNOWLEDGEMENT

This research is ostensibly supported by DIPA PNPB, Faculty of Science and Technology and Eijkman Biomolecular Institute.

REFERENCES

- [1] Baumgartener, T.G., Van Way, C.W., Vitamins in Nutrition Secrets, 26 adephia, Hanley and Belfus, 1999.
- [2] Borgna-Pignatti, C., Galanello, R., *Wintrobe's Clinical Hematology* 23 9, 11, 1319-1365.
- [3] De Franceschi, L., Cappellini, M.D., Olivieri, O., *Seminars in Thrombosis and Hemostasis* 2011, 37, 226-236.
- [4] Dhawan, V., Kumar, Kh. R., Marwaha, R.K., Naravan, S., Kamgar, M., *Ind. Pedi.* 2005, 42, 1141-1145.
- [5] Harborne, J.B., *Phytochemical Methods Guides Modern Way of Analyzing Plants*. Institut Teknologi Bandung, 1987.
- [6] Herawati, N., Jalaluddin, N., La Daha, Zenta, F., *Majalah Farmasi* 14 *Farmakologi* 2011, 15(1), 23-25.
- [7] Kang J., Li, Z., Wu, T., Jensen, G.S., Schauss, A.G., *Food Chemistry* 2010, 122, 3.
- [8] Latief, M., Nazaruddin and Nelson, *Prosiding SEMIRATA 2015 bidang MIPA BKS-PTN Barat Universitas Tanjungpura* 2015, 112 - 12.
- [9] Livingstone, F.B., *Frequencies of hemoglobin variants*. New York 24 Oxford: Oxford University Press, 1985.
- [10] Luqman, S., S.I., *Phytother Res* 2006, 20(4), 303-6.
- [11] Mahdi, E.A., *Acta Chim. Pharm. Indica* 2014, 4(3), 137-145.
- [12] Milena, R., Branka, Z., Biljana, S., Belton, N., Canat, B., *Thalassaemia Syndrome in Serbia, Hemoglobin* 2010, 34(5), 477-15.
- [13] Modell, B., D. 22 on, M., Birgens, H., Cario, H., Faustino, P., Giordano, P., *Scand J Clin Lab Invest* 2007, 67, 39-69.
- [14] Modell, B., *Bulletin of the World Health Organization* 2008, 86(6), 11-487.
- [15] Simsek, F., Ozturk, G., Kemahl, S., *Oxidant and Antioxidant Status* 1 *Beta Thalassemia Major Patients* 2008, 58, 34-38.
- [16] Thein, S.L., *Molecular basis of β -thalassemia*. Cold Spring Harb Perspect Med, 20 7.
- [17] Weatherall, D. J., *Annals of Human Biology* 2005, 32(2), 117-122.
- [18] Rajagukguk, R., Kosim, S., Tamam, M., *Medica Hospitalia* 2014, 27: 98-100.
- [19] Muhaimin, Syamsurizal, Chaerunisaa, A.Y., S 6 ga, M.S., *International Journal of ChemTech Research* 2016, 9(5), 418-424.
- [20] Muhaimin, Syamsurizal 6 Latief, M., Anggraini, S., Mujahidin, D., *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 2017, 8(1S), 141-148.

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

12%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1

www.mdpi.com

Internet Source

1%

2

www.cap-etest.com

Internet Source

1%

3

Lucia De Franceschi, Mariarita Bertoldi, Alessandro Matte, Sara Santos Franco et al. " Oxidative Stress and -Thalassemic Erythroid Cells behind the Molecular Defect ", Oxidative Medicine and Cellular Longevity, 2013

Publication

1%

4

Ngatijo, Rahmat Basuki, Bambang Rusdiarso, Nuryono. "Sorption-desorption profile of Au(III) onto silica modified quaternary amines (SMQA) in gold mining effluent", Journal of Environmental Chemical Engineering, 2020

Publication

1%

5

M Latief, A Utami, H Amanda, Muhaimin, Z Afifah. " Antioxidant activity of isolated compound from perepat roots () ", Journal of Physics: Conference Series, 2019

1%

6	Submitted to iGroup Student Paper	1 %
7	www.intechopen.com Internet Source	1 %
8	E. Diaz- Mesa, A. Fumero-Morera, P. Abreu-Gonzalez, E. Hernandez-Padron et al. "P.871 Relationship between therapeutic compliance with antipsychotics and total antioxidant status in acute paranoid schizophrenia", European Neuropsychopharmacology, 2020 Publication	1 %
9	www.scielo.org Internet Source	1 %
10	www.tandfonline.com Internet Source	1 %
11	Zahraa Falah Abd Alaali, Najat Mohammed Flyyih, Radhia hussain fadel. "Study of some clinico-pathological parameters of Thalassemia patients in AL-Zahraa hospital", Journal of Physics: Conference Series, 2019 Publication	1 %
12	Submitted to Trinity College Dublin Student Paper	1 %

13	Submitted to Lebanese American University Student Paper	1 %
14	d-nb.info Internet Source	1 %
15	Sezaneh Haghpanah, Nader Cohan, Mohammadreza Bordbar, Asghar Bazrafshan et al. "Effects of three months of treatment with vitamin E and N-acetyl cysteine on the oxidative balance in patients with transfusion-dependent β -thalassemia", Annals of Hematology, 2020 Publication	1 %
16	docplayer.net Internet Source	1 %
17	haematologica.org Internet Source	1 %
18	www.coursehero.com Internet Source	1 %
19	Submitted to King's College Student Paper	1 %
20	repository.unp.ac.id Internet Source	1 %
21	Jung Hoon Kang, Won Sik Eum. "Enhanced oxidative damage by the familial amyotrophic lateral sclerosis-associated Cu,Zn-superoxide dismutase mutants", Biochimica et Biophysica	<1 %

22	banglajol.info Internet Source	<1 %
23	onlinelibrary.wiley.com Internet Source	<1 %
24	Rajapakse, N.. "Inhibition of free radical-mediated oxidation of cellular biomolecules by carboxylated chitooligosaccharides", Bioorganic & Medicinal Chemistry, 20070115 Publication	<1 %
25	e-dergi.marmara.edu.tr Internet Source	<1 %
26	link.springer.com Internet Source	<1 %
27	www.eurekaselect.com Internet Source	<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

Artikel

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
