

Budiati

by Budiati Budiati

Submission date: 02-May-2023 07:50PM (UTC+0700)

Submission ID: 2081996302

File name: 1111budiati.pdf (203.54K)

Word count: 6178

Character count: 33743

IMPROVED TOLERANCE OF CHILI AGAINST DROUGHT STRESS BY THE USE OF *TRICHODERMA*-BASED COMPOST AND LIQUID FERTILIZER FROM LOCAL MICROORGANISM (LMO)

Eliyanti ELIYANTI*, Zulkarnain ZULKARNAIN*, Budiyati ICHWAN*, Adriani ADRIANI**

*⁴⁶ University of Jambi, Faculty of Agriculture, Department of Agroecotechnology, Ka⁴³ Pinang Masak, Jambi, Indonesia 36361

**⁴⁶ University of Jambi, Faculty of Animal Husbandry, Department of Animal Science, Kampus Pinang Masak, Jambi, Indonesia 36361

Corresponding author: Zulkarnain Zulkarnain, University of Jambi, Faculty of Agriculture, Department of Agroecotechnology, Kampus Pinang Masak, Jl. Raya Jambi – Muara Bulian KM 15, Jambi, Indonesia, phone: +628127864597, email: dr.zulkarnain@yahoo.com

Abstract. The present study was aimed to evaluate the effects of six different technology packages to improve tolerance of three chili cultivars for drought stress conditions. Six technology packages namely trichocompost and LMO spray every 7 days, trichocompost and LMO spray every 14 days, trichocompost only, LMO spray every 7 days only, LMO spray every 14 days only, and a control (no trichocompost nor LMO spray) were applied on three chili cultivars (Lado, Laris and Vitra). The trial was arranged in a randomized block design with three replicates consisting of four individual plants. Results indicated that the enrichment of growing media with trichocompost along with spraying of LMO improved the growth and yield of Lado, Laris and Vitra cultivars in drought conditions. Trichocompost in combination with LMO spray at 7-day interval increased the number of productive branches, as well as fruit number and fruit weight up to 28%, 58.84% and 64.28% in the cultivar Vitra, which was higher than the Laris and Lado. The application of *Trichoderma*-based compost plus spraying LMO every 7 days was able to increase tolerant properties of the three chili cultivars for drought stress. Within cultivars, Vitra was proven to be more adaptive and more tolerant against drought stress in comparison with Laris and Lado cultivars.

Keywords: *Capsicum annuum*; biofertilizer; abiotic stress; *Pomacea canaliculata*.

INTRODUCTION

Climate change occurring around the world has caused drought threat in many areas, and plants will suffer from a water scarcity. Drought is an important threat to sustainable agriculture because most of plant physiological processes are water-dependent [36]. Water shortage due to drought will inhibit nutrient uptake causing nutrient deficiency and disrupting cellular metabolism. This will lead to poor plant growth and development, resulting serious loss of yield [1, 26].

In tropical region such as Indonesia, limited soil water availability normally took place during dry season. Therefore, chili (*Capsicum annuum* L.) as many other vegetable crops, grown during dry season will experience drought stress unless water is adequately supplied by any method of irrigation. Ichwan et al. [14] reported that chili grown on soil with 50% field capacity showed slow growth and reduced yield. Also, number of productive branches, number of fruits and total fruit weight per plant were also low.

Drought is a condition of prolonged restricted ground water availability as a result of unfavorable conditions of growing medium. Many metabolism processes of plant were affected by drought stress, either directly or indirectly [38]. Noroozlo et al. [26] suggested that a significant change in cellular metabolism was generally occurred resulting in reduced plant growth and production. In addition to affecting plant physiological processes, drought also affects biochemical processes, anatomy and plant morphology [2].

Efforts to improve soil water availability during limited water supply can be done by the use of organic materials such as *Trichoderma*-enriched compost or trichocompost [6, 18, 22]. The use of organic materials may increase plant tolerance to drought due to better

soil structure, increase aeration in root zone, reduce mass density, increase cation exchange capacity, and maintain primary nutrients such as N and P [42]. The application of organic materials in soil has also proven to be beneficial to increase soil water holding capacity. Bhadha et al. [7] reported that an increase of 1% soil organic matter significantly increased soil water holding capacity.

The application of beneficial microorganisms such as *Trichoderma* in growing media [58] another way to enhance organic matter and improve chemical, physical and biological properties of soil. Microorganisms in compost are generally saprophyte bacteria (chemoorganotrophic and chemolithotrophic groups) and fungi. Many reports on the application of *Trichoderma*-enriched compost to promote plant growth and yield have been found in various horticultural crops such as *Lycopersicon esculentum* [24], *Brassica oleracea* [32] and *Ipomoea reptans* [43].

Bacteria from compost are mostly decomposers, also known as saprophytes, belonging to the group of chemoorganotrophic microorganisms. Among them, bacterial genera like *Pseudomonas*, *Burkholderia*, *Zymomonas*, *Xanthomonas*, and nitrogen-fixing aerobic bacteria can be found. Chemolithotrophic microorganisms can also be found in the compost like nitrifying bacteria that convert the ammonium into nitrites and nitrates. The most representative genera of this type are *Nitrosomonas* and *Nitrospira* [16, 20, 34, 35]. In addition, Fungi are also decomposers in nature and they can be found during the first and final phase of the composting process. The most representative genera are *Aspergillus*, *Acremonium*, *Chrysosporium*, *Fusarium*, *Mortierella*, *Penicillium*, and *Trichoderma* [3, 29].

Trichoderma was found to have antagonistic potential against root pathogenic isolates, so that it can

inhibit or even kill pathogenic fungi and increase plant tolerance to diseases caused by soil-borne pathogens [8, 9, 25, 41]. Therefore, trichocompost can be used as an alternative of organic fertilizers to increase plant growth and production while increasing crop protection.

Applying a liquid of local microorganisms (LMO) through foliar spray could increase the performance of plants grown on *Trichoderma*-enriched media as reported by many authors [10, 32, 43]. The LMO is a liquid containing various microorganisms involved in the overhauling of various organic wastes [27]. The main ingredients of LMO are carbohydrates, glucose, and a source of beneficial microorganisms, so they are recommended as decomposers, organic fertilizers, or biological pesticides. The ability of microorganisms in LMO to decomposition has been widely used for activation in the composting process [5, 21, 27].

One of potential source of ingredients of LMO is golden apple snail (*Pomacea canaliculata* Lamarck). It is a freshwater mollusk which is an important pest for rice cultivation in Indonesia. However, this snail is also able to provide benefits for farmers by processing it into LMO ingredients. LMO which is made from golden apple snails contains growth hormones that stimulate plant growth. In addition [27], LMO made of golden apple snails also contains nitrogen-fixing bacteria such as *Azospirillum*, *Azotobacter* [60] *Pseudomonas*, phosphate solubilizing microbes as well as microbes that play a role in the process of breaking down macromolecules into basic materials, various enzymes and growth hormones that are useful for plants [4, 17, 30].

The use of *Trichoderma* [56] the decomposition of organic matter may accelerate the composting process, as well as improve plant root system, and thereby increasing the ability of plants to extract water from soil. When combined with the application of LMO, it is believed that the technology will be able to overcome the lack of nutrient availability while increasing plant tolerance to drought stress. To our knowledge, there has no simultaneous application of trichocompost and LMO golden apple snail as a technology package in organic cultivation of chili was found, especially in the chili production centers in Jambi Province, Indonesia. Therefore, the use of this technology package could be a strategic step to support organic chili production in Jambi Province, and is worthy of further investigation.

MATERIAL AND METHODS

Experimental design and treatments

The study was conducted at the Teaching and Research Farm Faculty of Agriculture University of Jambi, Indonesia [57] from April 2020 through to September 2020. A factorial randomized block design with three replicates was employed in the trial. The variables tested were six technology packages (trichocompost and LMO spray every 7 days, trichocompost and LMO spray every 14 days,

trichocompost only, LMO spray every 7 days only, LMO spray every 14 days only, and no trichocompost nor LMO spray as control) that were applied on three chili cultivars (Lado, Laris and Vitra). Therefore, there were 18 treatment combinations.

LMO and trichocompost preparation

The LMO is fermented liquid made from natural ingredients containing microorganisms with the potential to transform organic matter, stimulate plant growth, and control pests and plant diseases. In this study, the mixture of LMO was obtained from golden apple snail, a fresh water mollusk, which is an important pest for rice cultivation in Indonesia. Some non-pathogenic microorganisms contained in a LMO solution are *Bacillus* sp., [27] *Sacharomyces* sp., *Azospirillum* sp., *Azotobacter* sp., *Pseudomonas* sp., *Aspergillus* sp. and *Lactobacillus* sp.

The LMO was made by preparing 5 kg crushed snails and 1 kg brown sugar into 10 L coconut water, stirred and mixed thoroughly. The mixed material was kept in a closed container connected to a small bottle filled with water through a small hose to neutralize gaseous materials formed due to fermentation, then left for approximately 2 weeks before use. A well-prepared LMO solution is characterized by the absence of a foul odor, and is ready to be applied to plants.

This study used *Trichoderma harzianum* isolated from soil samples which previously were used to grow chili. The isolation and propagation protocol followed Wagner [39] method. One [53] μm of soil sample from root area was mixed with 9 mL of sterile water in a test tube and stirred evenly. A serial dilutions were carried out so that the population could be calculated. One [17] mliter of the soil sample suspension was mixed with Potato Dextrose Agar (PDA) medium, and incubated at room temperature. The identification and morphological characterization of *Trichoderma* was determined using determination key of Watanabe [40] until the isolate was identified. [17] isolates of *T. harzianum* were then purified in PDA medium, and incubated for 7 days at room temperature. The whole series of work was done aseptically.

Trichocompost materials were prepared by mixing vegetable wastes, husk charcoal and cow manure in a definite proportion. Dolomite lime was added to maintain the pH neutral. A solution of brown sugar plus Effective Microorganism-4 (EM-4) were drenched to the materials, mixed thoroughly, and leave fermented for one week. After one week of fermentation, *T. harzianum* isolates were added to the materials, completely mixed, and keep fermented for next three weeks. Following three weeks of fermentation the *Trichoderma* fungus must have grown indicated by the appearance of fine white hyphae on the surface of materials, and this is the trichocompost which is ready to use as biofertilizer.

Plant preparation

Seeds were soaked in LMo solution for approximately 1 hour to induce uniform germination. Seeds were sown on a mixture of soil + trichocompost + sand (2:1:1) in a plastic pot. Spacing up to two seeds per pot was done at 2 weeks after germination, and following 4 weeks of germination the seeds were ready to be transplanted to black polyethylene bags (polybags) in greenhouse.

The growing medium in the greenhouse was 10 kg mixture of soil and trichocompost with a ratio of 1:1. The LMo was sprayed at 500 mL per plant every 7 or 14 days according to treatment. Routine plant maintenance such as pest and disease control, weeding and water control were carried out in order to keep plants in drought conditions.

Plants were subjected to drought from the beginning of transfer to polybags in the greenhouse (0 day after transplanting). The drought condition was made by keeping soil water content at 75% of field capacity, which was obtained through gravimetric measurements [37]. This was done by pouring water until reach saturation, and water was let to drain for two days. The difference between initial weight of the polybags and final weight after draining indicated the amount of water given to soil to reach field capacity condition. If the amount of water was 100 mL to reach field capacity, so the amount of water to reach 75% of field capacity would be 75 mL. The volume of water was applied to the plant every two days to keep drought condition.

Data collected and statistical analysis

Observations and data collected were made on the following variables: plant height (cm), number of productive branches, days to flowering, number of fruits per plant, weight of fruit per plant (kg), and relative water content. Except for RLWC, data on the effect of trichocompost treatment and LMo spraying were statistically analyzed using Analysis of Variance, followed by Duncan's Multiple Range Test to see the difference in the effect among treatment means.

Plant height was measured at 12 weeks after transplanting from the root neck to the tip of the stem; number of productive branches was measured when dichotomous began to appear until 12 weeks after transplanting; days to flowering was measured when the first flower appeared on the sample plants; number of fruits per plant was measured by adding up all fruits obtained from the first to the last harvest; weight of fruit per plant was measured by weighing all fruits yielded from the first to the last harvest.

Table 1. Summary of analysis of variance (ANOVA) the effect of technology packages (trichocompost + LMo) and cultivars on growth and production variables of chili grown under drought condition (** = highly significant; * = significant; ns = non-significant)

Source of Variations	Plant height (cm)	Number of productive branch	Days to flowering	Number of fruit	Weight of fruit (g)
Block	**	ns	*	ns	ns
Plant Cultivars (C)	**	ns	ns	**	**
Technology Package (T)	**	**	ns	**	**
Interaction (T × C)	ns	ns	ns	**	**
Coefficient of Variance (%)	5.86	10.07	22.35	6.44	7.29

20

Relative leaf water content (RLWC) was determined according to González and González-Vilar [11] method. Young leaf samples were taken randomly from 3 sample plants of 3 replicates at 10 weeks after transplanting. Six sections of leaves (apical and basal parts were removed) from a same plant were put in a pre-weight tube, sealed and placed in 10°C to prevent growth and evaporation. Tubes with tissue sections inside were weight to give a value for tissue fresh weight (FW). Once every sample has been weighed, a small volume of distilled water was added. Tubes were placed a fridge for 24 hours to allow the tissue taking up water. The tissue sections were taken away from distilled water, the excess of water on the leaf surface were removed carefully with tissue paper, and put in a pre-weight tube to give a measure of fully turgid fresh weight (TFW). The opened-tubes were then placed in an drying oven (70°C) for 24 hours when the tissues were reweigh to obtain a value for dry weight (DW). Relative leaf water content (RLWC) was then calculated using the following equation:

$$RLWC = \frac{FW - DW}{TFW - DW} \times 100$$

RESULTS

Data on experimental variables were statistically analyzed using analysis of variance (ANOVA) to detect the effect of different cultivars and technology packages on growth and production of chili. The results of the ANOVA is summarized in Table 1.

Plant height

The ANOVA presented in Table 1 indicates that both cultivars and technology packages significantly affect plant height, but the interaction of these two factors does not show any significant effect on plant height. The effect of different cultivars and technology packages on plant height is presented in Table 2 and Table 3.

Table 2 shows that Laris and Vitra significantly grow higher than Lado in drought conditions. This indicates that Laris and Vitra are more adaptable to sub-optimal or drought conditions than Lado. In drought conditions, the presence of *Trichoderma*-based compost in the growing medium and spraying LMo at intervals of 7 or 14 days were able to increase the growth of chili plants by 15% (96.16 cm) and 8% (90.60 cm) respectively, compared to without compost nor *Trichoderma* microorganisms (83.68 cm) as indicated in Table 3.

Productive branches

The ANOVA presented in Table 1 shows that cultivars do not significantly affect the number of productive branches. In contrast, the technology packages show a significant effect on the number of productive branches. However, the interaction between these two variables do not show any significant effect on the number of productive branches. The effect of technology packages on the number of productive branches is presented in Table 4.

Table 4 shows that under drought conditions, growing media enriched with trichocompost is able to increase the number of productive branches by 29% (118.76) to 38% (127.30) compared to control (92.03) either with or without spraying of LMO.

Days to flowering

Days to flowering was recorded at the earliest time of floral development and flower buds can be seen with bare eyes on leaf axil. The ANOVA presented in Table 1 shows that neither cultivars nor technology

packages nor interaction of these two variables significantly affect days to flowering.

Number of fruits

The ANOVA presented in Table 1 indicates that either technology packages or cultivars or interaction of these two variables significantly affect chili production in term of number of fruit. The effect of different cultivars and technology packages as well as the interaction of the two variables on the number of fruit production is presented in Table 5.

Table 5 shows that enriching growing medium with trichocompost and spraying LMO could increase the average number of fruits per plant under drought conditions. In general, Vitra shows better response than Lado and Laris cultivars. The highest average number of fruit per plant is shown by Vitra grown on medium enriched with trichocompost and sprayed with LMO at 7-day interval (87.67), followed by 14-day interval (82.00).

Table 2. The effect of different cultivars on the height of chili plants grown under 75% field capacity

Chili cultivars	Average plant height (cm)
Laris	96.68 a
Vitra	96.31 a
Lado	77.11 b

Numbers followed by the same lowercase indicate a non-significant different based on Duncan's Multiple Range Test at $\alpha = 0.05$.

Table 3. The effect of different technology packages on the height of chili plants grown under 75% field capacity

Technology packages	Average plant height (cm)
Trichocompost and LMO spray at 7-day interval	96.16 a
Trichocompost and LMO spray at 14-day interval	90.60 b
LMO spray at 7-day interval	90.37 b
Trichocompost only	89.87 b
LMO spray at 14-day interval	89.52 b
Control	83.68 c

Numbers followed by the same lowercase indicate a non-significant different based on Duncan's Multiple Range Test at $\alpha = 0.05$.

Table 4. The effect of different technology packages on the number of productive branches in three chili cultivars grown under 75% field capacity

Technology packages	Average productive branch
Trichocompost and LMO spray at 14-day interval	127.30 a
Trichocompost only	120.63 a
Trichocompost and LMO spray at 7-day interval	118.76 a
LMO spray at 7-day interval	101.99 b
LMO spray at 14-day interval	99.36 b
Control	92.03 b

Numbers followed by the lowercase indicate a non-significant different based on Duncan's Multiple Range Test at $\alpha = 0.05$.

Table 5. The effect of different technology packages and different cultivars on fruit number of chili grown under 75% field capacity

Technology packages	Chili cultivars		
	Lado	Laris	Vitra
Control	26.00 c	32.33 b	35.67 a
	F	F	E
LMO spray at 7-day interval	45.33 b	42.67 c	50.00 a
	E	E	D
LMO spray at 14-day interval	52.00 a	47.00 c	48.00 b
	D	D	E
Trichocompost only	67.67 b	60.67 c	74.00 a
	C	C	C
Trichocompost and LMO spray at 7-day interval	79.33 b	67.67 c	87.67 a
	A	A	A
Trichocompost and LMO spray at 14-day interval	73.67 b	63.33 c	82.00 a
	B	B	B

Numbers followed by the same lowercase in rows and the same uppercase in columns indicate non-significant different based on Duncan's Multiple Range Test at $\alpha = 0.05$.

Table 6. The effect of different technology packages and different cultivars on fruit weight (g) of chili grown under 75% field capacity

Technology packages	Chili Varieties		
	Lado	Laris	Vitra
Control	79.67 b F	99.17 a F	114.03 a E
LMo spray at 7-day interval	145.53 b E	142.27 b E	170.83 a C
LMo spray at 14-day interval	170.30 a D	152.25 b D	150.65 b D
Trichocompost only	233.17 b C	191.67 c C	267.25 a C
Trichocompost and LMo spray at 7-day interval	303.43 b A	248.55 c A	318.26 a A
Trichocompost and LMo spray at 14-day interval	295.13 b B	231.32 c B	305.68 a B

Numbers followed by the same lowercase in rows and the same uppercase in columns indicate non-significant different based on Duncan's Multiple Range Test at $\alpha = 0.05$.

Table 7. The effect of different technology packages and different cultivars on relative leaf water content (%) of chili grown under 75% field capacity

Technology packages	Chili Varieties		
	Lado	Laris	Vitra
Trichocompost and LMo spray at 14-day interval	71.81 ± 1.905	79.17 ± 1.801	86.44 ± 1.183
Trichocompost and LMo spray at 7-day interval	67.91 ± 1.238	72.02 ± 1.799	84.23 ± 2.170
Trichocompost only	62.97 ± 1.158	63.32 ± 0.400	77.20 ± 1.004
LMo spray at 7-day interval	61.57 ± 1.697	57.44 ± 1.706	78.69 ± 1.198
LMo spray at 14-day interval	61.23 ± 1.132	55.04 ± 3.290	76.00 ± 2.797
Control	57.72 ± 1.401	54.25 ± 3.614	61.70 ± 1.053

Relative leaf water contents was determined on individual leaves taken from 3 sample plants of 3 replicates at 10 weeks after transplanting.
± Standard Error

Weight of fruits

The ANOVA presented in Table 1 indicates that either technology packages or cultivars or interaction of these two variables significantly affect chili fruit weight. The effect of different technology packages and cultivars as well as the interaction of the two variables on the average fruit weight is presented in Table 6.

Similar to the average number of fruits per plant, enrichment of growing medium with *Trichoderma*-based compost and spraying LMo also increased average fruit weight per plant as presented in Table 6. The greatest fruit weight (318.26 g) was obtained by Vitra cultivar which was planted on medium enriched with *Trichoderma*-based compost and sprayed with LMo every 7-day, followed by spraying every 14-day intervals (305.68 g) on the same medium.

Relative leaf water content

Data presented in Table 7 indicate that under limited soil water availability there is variation in relative leaf water content (RLWC) among different technology packages and chili cultivars. The cultivar Vitra has a higher RLWC (77.38%) than Lado (63.87%) and Laris (63.54%). A higher RLWC indicates a quick plant physiological response against extreme environmental conditions such as drought. All chili cultivars show a higher RLWC when they are grown on media enriched with *Trichoderma*-based compost and sprayed with LMo at either 7- or 14-day intervals (79.14% and 74.72%, respectively). This indicates that the availability of *Trichoderma*-based compost and the application of LMo could increase physiological response of plants grown under drought condition.

DISCUSSION

Many reports indicated that soil water availability during drought could be improved by the use of *Trichoderma*-enriched compost or trichocompost [6, 18, 22]. In our study, the application of trichocompost along with LMo technology packages had a significant effect on plant height, number of productive branches, number of fruits and fruit weight of the three chili cultivars grown under drought conditions. There was also a significant effect of the interaction between the cultivars and LMo technology packages on fruit number and fruit weight (Table 1). However, neither cultivars nor technology packages nor their interactions showed significant effect on flowering. Days to flowering of the three chili cultivars ranged from 26 to 28 days after planting. In normal conditions and based on the descriptions of the cultivars, the average days to flowering of Lado, Laris and Vitra were 25 to 30 days after planting. This indicates that drought did not affect floral growth and development of the three chili cultivars when trichocompost and LMo were applied. However, Mitra [23] stated that days to flowering was dependent on altitude and genetic characteristics. This is in accordance with reports of Lembang Vegetable Research Institute [19] that days to flowering of chili cultivars grown in lowlands was approximately 28 days after transplanting and in uplands ranging from 34 to 39 days after transplanting.

This study revealed that the cultivar Vitra was more tolerant to drought stress than Laris and Lado. This was indicated by higher fruit number and fruit weight which were nearly as much as normal growing conditions. On the other hand, the Lado produced lower fruit number and fruit weight than Laris and Vitra (Tables 5 and 6). This means that Lado is a less tolerant cultivar against drought stress, while Laris is moderately tolerant.

The application of trichocompost along with LMO spray at 7-day interval was able to optimize the development of productive branches and fruit formation in the three chili cultivars grown under drought conditions. This is in accordance with the finding of Hasyim et al. [12] and Rizki et al. [31] who reported that trichocompost could increase the weight of fruit per plant by more than 70% in chili.

Genotype is an important factor in building plant tolerance to drought stress. In line with this, Rosmaina et al. [33] studied the indicators of drought stress tolerance based on correlation and cross-analysis of normal condition (100% field capacity) and stress conditions (75%, 50% and 25% field capacity). It was found that the number of fruit had a positive correlation and had a direct effect on fruit weight per plant, so that it became a phenotypic character that could be used as a selection criterion for drought tolerant chilies. Thus, cultivars that are able to grow and produce higher fruit number and fruit weight under drought conditions can be categorized as drought stress tolerant cultivars.

The application of trichocompost-enriched media along with LMO spray at 7-day interval was able to increase the number of fruit by up to 58.84% and fruit weight by up to 64.28% in Vitra compared to control. This yield was higher than those achieved in Lado and Laris cultivars. This also indicates that the use of trichocompost coupled with LMO spray every 7 days could reduce the adverse effect of drought stress in the three cultivars, especially in Vitra. This is in accordance with the findings in previous study that foliar spray of water soluble nutrients and biostimulants was proven to increase the ability of the leaves to maintain turgor pressure allowing the nutrients contained in the solution to enter the stomata [13, 14, 15].

In drought stress conditions, Vitra was found to be more tolerant than Laris or Lado. This indicated by higher number of productive branches, number of fruit, and fruit weight. This tolerance response was also reinforced by a high relative leaf water content (RLWC) (nearly 80%) or close to stress-free conditions, indicating that Vitra had better drought stress tolerance than the other two genotypes. This was in line with Okunlola et al. [28] report that drought stress conditions significantly reduced RLWC values in *C. chinense*, *C. annum* and *C. frutescense*. Plants grown under normal condition had an RLWC of 93.75% - 95.05%, plants under light drought stress had an RLWC of 90.60% - 93.40%, plants under moderate drought stress had an RLWC of 69.30% - 80.90%, and plants under severe drought stress had an RLWC of 53.25% - 68.95%.

In conclusion, our study showed that the enrichment of growing media with *Trichoderma*-based compost (trichocompost) along with spraying local microorganisms (LMO) were important to improve tolerant properties of three chili cultivars (Lado, Laris and Vitra) for drought stress. Under drought stress condition, trichocompost and routine spray of LMO increased the number of productive branches, number of fruits and weight of fruits by 28%, 58.84% and 64.28% respectively in Vitra, which was higher than Laris and Lado. This indicates that Vitra is a more

adaptive and more tolerant cultivar than Laris and Lado cultivars for drought stress.

Acknowledgements. The authors would like to thank Rector University of Jambi for financial support provided through Internal Competitive Research Grant (Contract No. 572/50/21.18/PG/SPK/2020, dated 20th April 2020).

Conflict of interest. There is no actual or potential conflict of interest in relation to this article.

REFERENCES

- Ahmadi, M., Souri, M.K., (2018): Growth and mineral elements of coriander (*Coriander sativum* L.) plants under mild salinity with different salts. *Acta Physiologica* 41: 166-173.
- Ai, N.S., Banyo, Y., (2011): Konsentrasi klorofil sebagai indikator kekurangan air pada tanaman. *Jurnal Ilmiah Sains*, 33: 166-173.
- Anastasi, A., Varese, G., Marchisio, V., (2005): Isolation and identification of fungal communities in compost and micompost. *Mycologia*, 97: 33-44.
- Artyszak, A., Gozdowski, D., (2020): The effect of growth activators and Plant Growth-Promoting Rhizobacteria (PGPR) on the soil properties, root yield, and technological quality of sugar beet. *Agronomy*, 10: 1262.
- Astuti, T., Amir, Y.S., Yelni, G., Isyaturriyadhah, I., (2014): The result of biotechnology by local microorganisms to banana peel on rumen fluid characteristics as ruminant feed. *Journal of Advanced Agricultural Technologies*, 1: 28-31.
- Bae, H., Sicher, R.C., Kim, M.S., Kim, S.-H., Strem, M.D., Melnick, R.L., Bailey, B.A., (2009): The beneficial endophyte *Trichoderma hamatum* isolate DIS 219b promotes growth and delays the onset of the drought response in *Theobroma cacao*. *Journal of Experimental Botany*, 60: 3279-3295.
- Bhadha, J.H., Capasso, J.M., Khatiwada, R., Swanson, S., LaBorde, C., (2017): Raising soil organic matter content to improve water holding capacity. IFAS Extension University Florida, Gainesville, Florida, pp. 5.
- Brotman, Y., Lisek, J., Meret, M., Chet, I., Willmitzer, L., Viterbo, A., (2012): Transcript and metabolite analysis of the *Trichoderma*-induced systemic resistance response to *Pseudomonas syringae* in *Arabidopsis thaliana*. *Microbiology*, 158: 139-146.
- Contreras-Cornejo, H.A., Macías-Rodríguez, L., Beltrán-Pena, E., Herrera-Estrella, A., López-Bucio, J., (2011): *Trichoderma*-induced plant immunity likely involves both hormonal- and camalexin independent mechanisms in *Arabidopsis thaliana* and confers resistance against necrotrophic fungus *Botrytis cinerea*. *Plant Signaling and Behavior*, 6: 1445-1563.
- dos Santos, J., Gomes, D.C.B.B., Nabais, C.N., (2018): The influence of local microorganism liquid fertilizer concentration from banana stumps and NPK fertilizer dosage against kangkung or kale growth (*Ipomoea reptans*), conducted at NCBA. *International Journal of Development Research*, 8: 24626-24634.
- González, L., González-Vilar, M., (2001): Determination of Relative Water Content, pp. 207-212. In Roger, M.J.R. [ed.]: *Handbook of Plant Ecophysiology Techniques*. Kluwer Academic Publishers, Netherlands.
- Hasyim, A., Setiawati, W., Lukman, L., (2015): Technological innovation of sustainable pest and disease management on chili peppers: an alternative effort to establish harmonious ecosystems. *Pengembangan Inovasi Pertanian*, 8: 1-10.
- Ichwan, B., Eliyanti, E., Zulkarnain, Z., (2021): Effect of biostimulants and media compositions on growth and yield of *Capsicum annum* L. under drought stress conditions. *Advances in Horticultural Science*, 32: 129-137.

- 26
- [14] Ichwan, B., Suwignyo, R.A., Susilawati, (2017): Response of red chilli varieties under drought stress. *Russian Journal of Agricultural and Socio-Economic Sciences*, 6: 361-368.
- 63
- [15] Ichwan, B., Suwignyo, R.A., Susilawati, S., Eliyanti, E., Zulkarnain, Z., (2021): Foliar spray of water soluble fertilizer enhances drought tolerance of chili pepper. *Analele Universității din Oradea, Fascicula Biologie*, 28: 19-24.
- [16] Insam, H., de Bertoldi, M., (2007): Microbiology of the composting process, pp. 25-48. In Diaz, L.F., de Bertoldi, M., Bidlingmaier, W., Goluke, C. [eds.]: *Compost Science & Technology*. Elsevier, Oxford, UK.
- [17] Kang, B.G., Kim, W.T., Yun, H.S., Chang, S.C., (2010): Use of plant growth-promoting rhizobacteria to control stress responses of plant roots. *Plant Biotechnology Reports*, 4: 179-183.
- [18] Khoshmanzar, E., Aliasgharzad, N., Neyshabouri, M.R., Khoshru, B., Arzanlou, M., Lajayer, B.A., (2020): Effects of *Trichoderma* isolates on tomato growth and inducing its tolerance to water-deficit stress. *International Journal of Environmental Science and Technology*, 17: 869-878.
- [19] Lembang Vegetable Research Institute, (2015): Guidelines control of virus diseases in chili plants. Horticulture Research and Development Center, Lembang Agricultural Research and Development Center, Bandung (in Indonesian), pp. 22.
- [20] Madigan, M.T., Bender, K.S., Buckley, D.H., Sattley, W.M., Stahl, D.A., (2021): *Biology of microorganisms* (8th edition). Pearson Education, Hoboken, NJ, pp. 1124.
- [21] Manullang, R.R., Rusmini, R., Daryono, D., (2018): Combination microorganism as local bio activator compost Kirinyuh. *International Journal of Scientific and Technology Research*, 7: 1-5.
- [22] Mastouri, F., Björkman, T., Harman, G.E., (2012): *Trichoderma harzianum* enhances antioxidant defense of tomato seedlings and resistance to water deficit. *Molecular Plant-Microbe Interactions*, 25: 1264-1271.
- [23] Mitra, J., (2001): Genetics and genetic improvement of drought resistance in crop plant. *Current Science*, 80: 758-763.
- [24] Molla, A.H., Haque, M.M., Haque, M.A., Ilias, G.N.M., (2012): *Trichoderma*-enriched biofertilizer enhances production and nutritional quality of tomato (*Lycopersicon esculentum* Mill.) and minimizes NPK fertilizer use. *Agricultural Research*, 1: 265-272.
- [25] Mukherjee, P.K., Horwitz, B.A., Kenerley, C.M., (2012): Secondary metabolism in *Trichoderma* - a genomic perspective. *Microbiology*, 158: 35-45.
- [26] Noroozlo, Y.A., Souri, M.K., Delshad, M., (2019): Stimulation effects of foliar applied glycine and glutamine amino acids on lettuce growth. *Open Agriculture*, 4: 164-172.
- [27] Nurhaita, N., Neli, D., Suliasih, S., (2015): Fermentation with local microorganism to improve pod cacao quality as ruminants feed. *Proceeding International Seminar on Promoting Local Resources for Food and Health*, 12-13 October 2015. Bengkulu, Indonesia: 218-221.
- [28] Okunlola, G.O., Olatunji, O.A., Akinwale, R.O., Tariq, A., Adelusie, A.A., (2017): Physiological response of the three most cultivated pepper species (*Capsicum* spp.) in Africa to drought stress imposed at three stages of growth and development. *Scientia Horticulturae*, 224: 198-205.
- [29] Pérez Bocourt, Y., Rebollido Ríos, R., Martínez Silva, J., (2010): Aislamiento e identificación de hongos en compost elaborado a partir de residuos sólidos urbanos (Isolation and identification of fungi in compost produced from municipal solid waste, in Spanish). *Agro Sur*, 38: 1-7.
- [30] Retnowati, Y., Katili, A.S., (2021): Identification of fermentative bacteria on local microorganisms of golden snail (*Pomacea canaliculata* Lamarck, 1822). *Biodiversitas*, 22: 778-784.
- [31] Rizki, H.B., Puspita, F., Adiwirman, (2015): Test some of the trichocompost formulated against growth and production of red chili. *Jurnal Online Mahasiswa*, 2: 1-14.
- [32] Roeswitawati, D., Ningsih, Y.U., Muhidin, M., (2018): The effect of local microorganism (MOL) concentration of banana hump and fruit waste on growth and yield of broccoli plants (*Brassica oleracea*). *Advances in Engineering Research*, 172: 310-314.
- [33] Samain, R., Sobir, S., Parjanto, P., Yunus, A., (2019): Correlations and path analysis of some characters in chili pepper (*Capsicum annum* L.) under normal and drought stress. *Jurnal Hortikultura*, 29: 147-158.
- [34] Ryckeboer, J., Mergaert, J., Coosemans, J., Deprins, K., Swings, J., (2003): Microbiological aspects of biowaste during composting in a monitored compost bin. *Journal of Applied Microbiology*, 94: 127-137.
- [35] Ryckeboer, J., Mergaert, J., Vaes, K., Klammer, S., De Clercq, D., Coosemans, J., Insam, H., Swings, J., (2003): A survey of bacteria and fungi occurring during composting and self-heating processes. *Annals of Microbiology*, 53: 399-410.
- [36] Seleiman, M.F., Al-Suhaibani, N., Ali, N., Akmal, M., Alotaibi, M., Refay, Y., Dindaroglu, T., Abdul-Wajid, H.H., Battaglia, M.L., (2021): Drought stress impacts on plants and different approaches to alleviate its adverse effects. *Plants*, 10: 259.
- [37] Shukla, A., Panchal, H., Mishra, M., Patel, P.R., Srivastava, H.S., Patel, P., Shukla, A.K., (2015): Soil moisture estimation using gravimetric technique and FDR probe technique: A comparative analysis. *American International Journal of Research in Formal, Applied and Natural Sciences*, 8: 89-92.
- [38] Souri, M.K., Hatamian, M., (2018): Aminocheleates in plant nutrition: a review. *Journal of Plant Nutrition*, 42: 67-78.
- [39] Wagner, J.E., (1993): *The antagonistic potency of several fungus species in cultivated soil in tropics*. Tata McGraw-Hill Publishing Co. Ltd, New Delhi, pp. 218.
- [40] Watanabe, T., (2010): *Pictorial atlas of soil and seed fungi: morphologies of cultured fungi and key to species (3rd edition)*. CRC Press, Boca Raton, pp. 426.
- [41] Widyastuti, S.M., Sumardi, S., Harjono, H., (1999): Potensi antagonistik tiga *Trichoderma* spp. terhadap delapan penyakit akar tanaman kehutanan. *Buletin Kehutanan*, 41: 32.
- [42] Williams, C.N., Uzo, J.O., Peregrine, W.T.H., (1991): *Vegetable production in the tropics*. Longman Scientific and Technical, Essex, England, pp. 179.
- [43] Yuliana, M., (2021): The effect of local microorganism (Mol) as liquid organic fertilizer to the growth of *Ipomea reptans* Poir. *Jurnal Biota*, 7: 51-56.

Received: December 21, 2021

Accepted: May 30, 2022

Published Online: June 2, 2022

Analele Universității din Oradea, Fascicula Biologie

<https://www.bioresearch.ro/revistaen.html>

Print-ISSN: 1224-5119

e-ISSN: 1844-7589

CD-ISSN: 1842-6433

University of Oradea Publishing House



ORIGINALITY REPORT

21 %
SIMILARITY INDEX

%
INTERNET SOURCES

21 %
PUBLICATIONS

%
STUDENT PAPERS

PRIMARY SOURCES

- 1** P. Umadevi, M. Anandaraj. "Proteomic analysis of the tripartite interaction between black pepper, *Trichoderma harzianum* and *Phytophthora capsici* provides insights into induced systemic resistance mediated by *Trichoderma* spp.", *European Journal of Plant Pathology*, 2019
Publication **1** %

- 2** Khalida Firda Zanatia, Cecep Hidayat, Esty Puri Utami. "Respons Tanaman Bawang Merah Terhadap Pemberian Pupuk Organik Cair Air Kelapa dan Mikroorganisme Lokal Bonggol Pisang", *Jurnal Pertanian Terpadu*, 2021
Publication **1** %

- 3** M. Szczałba, T. Kopta, M. Gąstoł, A. Sękara. "Comprehensive insight into arbuscular mycorrhizal fungi, spp. and plant multilevel interactions with emphasis on biostimulation of horticultural crops ", *Journal of Applied Microbiology*, 2019
Publication **1** %

4

R. Prajapati, D. Chakraborty, V. Kumar. "ADVANCES IN SOIL MOISTURE RETRIEVAL FROM NEAR-SURFACE MEASUREMENTS USING SATELLITE REMOTE SENSING", ISPRS - International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences, 2018

Publication

1 %

5

Hassan Etesami. "Potential advantage of rhizosheath microbiome, in contrast to rhizosphere microbiome, to improve drought tolerance in crops", Rhizosphere, 2021

Publication

1 %

6

Laily Ilman Widuri, Benyamin Lakitan, Jyunichi Sakagami, Shin Yabuta, Kartika Kartika, Erna Siaga. "Short-term drought exposure decelerated growth and photosynthetic activities in chili pepper (*Capsicum annum* L.)", Annals of Agricultural Sciences, 2020

Publication

1 %

7

Debmalya Dasgupta, Kulbhushan Kumar, Rashi Miglani, Rojita Mishra, Amrita Kumari Panda, Satpal Singh Bisht. "Microbial biofertilizers: Recent trends and future outlook", Elsevier BV, 2021

Publication

1 %

8

S. Klammer. "Bacterial community patterns and thermal analyses of composts of various

1 %

9

Atainah Ul Hukma, Meta Yuliana, Novin Teristiandi. "Pengaruh Pupuk Organik Cair (POC) Buah Belimbing Wuluh (*Averrhoa bilimbi* L.) terhadap Pertumbuhan Sawi Hijau", *EKOTONIA: Jurnal Penelitian Biologi, Botani, Zoologi dan Mikrobiologi*, 2022

Publication

1 %

10

Roberto N. Silva, Valdirene Neves Monteiro, Andrei Stecca Steindorff, Eriston Vieira Gomes et al. "Trichoderma/pathogen/plant interaction in pre-harvest food security", *Fungal Biology*, 2019

Publication

1 %

11

Mariana Petkova, Stefan Shilev. "Revealing Fungal Diversity in Mesophilic and Thermophilic Habitats of Sewage Sludge Composting by Next-Generation Sequencing", *Applied Sciences*, 2023

Publication

1 %

12

Salwa Latif, Amany Gaber Mohamed, Kuni Sueyoshi. "Effect of *Bacillus subtilis* on Some Physiological and Biochemical Processes in Barley (*Hordeum vulgare* L.) Plant Grown under Salt Stress", *Egyptian Journal of Botany*, 2020

Publication

1 %

13 Winih Sekaringtyas Ramadhani, Soemarno, Ali Rahmat, Priyo Cahyono, Fitri Wijayanti. "Efforts to reduce soil compaction in pineapple plantations using plants rotation with banana Cavendish in Central Lampung, Indonesia", AIP Publishing, 2023 1 %

Publication

14 Sandra Pérez-Álvarez, Eduardo Fidel Héctor Ardisana, Marco Antonio Magallanes-Tapia, Cesar Marcial Escobedo Bonilla et al. "Chapter 2 Microorganisms Used as Growth Regulators in Modern Agriculture", Springer Science and Business Media LLC, 2022 <1 %

Publication

15 I. P. Sapinas, L. K. Abbott. "Soil Fertility Management Based on Certified Organic Agriculture Standards – a Review", Sustainable Agriculture Research, 2020 <1 %

Publication

16 Khoshrooz Kazemi, Baiyu Zhang, Leonard M. Lye. "Assessment of Microbial Communities and Their Relationship with Enzymatic Activity during Composting", World Journal of Engineering and Technology, 2017 <1 %

Publication

17 Febri Doni, Che Radziah Che Mohd Zain, Anizan Isahak, F. Fathurrahman et al. "A simple, efficient, and farmer-friendly <1 %

Trichoderma-based biofertilizer evaluated with the SRI Rice Management System", Organic Agriculture, 2017

Publication

18

Manish Tiwari, Ritesh Kumar, Doohong Min, S V Krishna Jagadish. "Genetic and molecular mechanisms underlying root architecture and function under heat stress - A hidden story", Plant, Cell & Environment, 2022

Publication

19

Nuhaa Soobhany. "Insight into the recovery of nutrients from organic solid waste through biochemical conversion processes for fertilizer production: A review", Journal of Cleaner Production, 2019

Publication

20

Arantza del-Canto, Álvaro Sanz-Saez, Anna Sillero-Martínez, Eider Mintegi, Maite Lacuesta. "Selected indigenous drought tolerant rhizobium strains as promising biostimulants for common bean in Northern Spain", Frontiers in Plant Science, 2023

Publication

21

Boris Basile, Youssef Roupheal, Giuseppe Colla, Sebastian Soppelsa, Carlo Andreotti. "Appraisal of emerging crop management opportunities in fruit trees, grapevines and

<1 %

<1 %

<1 %

<1 %

berry crops facilitated by the application of biostimulants", *Scientia Horticulturae*, 2020

Publication

22

Cia, M.C., A.C.R. Guimarães, L.O. Medici, S.M. Chabregas, and R.A. Azevedo.

"Antioxidant responses to water deficit by drought-tolerant and -sensitive sugarcane varieties : Water-deficit responses by sugarcane", *Annals of Applied Biology*, 2012.

Publication

23

Nurhaita, Neli Definiati, Syahro Ali Akbar, Urip Santoso. "The effect of cocoa pod

fermentation with mol of rumen content on fiber fraction component and in vitro digestibility", *IOP Conference Series: Earth and Environmental Science*, 2019

Publication

24

Giacomo Bartoli, Stefania Bottega, Carmelina Spanò. "Morpho-anatomical and physiological traits of *Agrostis castellana* living in an active geothermal alteration field", *Biologia*, 2015

Publication

25

Lei Yan, Yu Zeng, Muhammad Riaz, Jin Cheng, Cuncang Jiang. "Exogenous proline triggered internal tolerance mechanism in trifoliate orange (*Poncirus trifoliata*) acclimated to boron-deficiency", *Scientia Horticulturae*, 2021

Publication

<1 %

<1 %

<1 %

<1 %

26

Tilen Zamljen, Vesna Zupanc, Ana Slatnar. "Influence of irrigation on yield and primary and secondary metabolites in two chilies species, *Capsicum annum* L. and *Capsicum chinense* Jacq", *Agricultural Water Management*, 2020

Publication

<1 %

27

"Microbiological Activity for Soil and Plant Health Management", *Springer Science and Business Media LLC*, 2021

Publication

<1 %

28

Solmaz Aslanzadeh, Katherine Kho, Irnayuli Sitepu. "An Evaluation of the Effect of Takakura and Effective Microorganisms (EM) as Bio Activators on the Final Compost Quality", *IOP Conference Series: Materials Science and Engineering*, 2020

Publication

<1 %

29

Shihomi Uzuhashi, Haruka Ikeda, Ai Kamekawa, Lorien E. Radmer et al. "Presence of two species-level groups in *Globisporangium splendens* isolates in Japan", *European Journal of Plant Pathology*, 2019

Publication

<1 %

30

Zainal Abidin, Rusmini Rusmini, Riama Rita Manullang, Daryono Daryono. "Kualitas Mikroorganisme Lokal Dari Keong Mas

<1 %

Dengan Berbagai Jumlah Bahan Yang berbeda", AGROSAINTIFIKA, 2022

Publication

31

Karin E. Limburg, Dennis P. Swaney, David L. Strayer. "River Ecosystems", Elsevier BV, 2022

Publication

<1 %

32

Nutrition & Food Science, Volume 34, Issue 3 (2006-09-19)

Publication

<1 %

33

Mara Gmez-Brandn, Marina Fernndez-Delgado, Jorge Domnguez, Heribert Insam. "Chapter 10 Animal Manures: Recycling and Management Technologies", IntechOpen, 2013

Publication

<1 %

34

S Hafsah, Yusnizar, Nura, K S Kaloko, F Reza, Firdaus. "Bio-priming of Aceh local chili seeds in the effort to increase production and begomo virus resistance", IOP Conference Series: Earth and Environmental Science, 2022

Publication

<1 %

35

André Alisson Rodrigues da Silva, Geovani Soares de Lima, Carlos Alberto Vieira de Azevedo, Hans Raj Gheyi et al. "Salicylic acid relieves the effect of saline stress on soursop morphology", Ciência e Agrotecnologia, 2021

Publication

<1 %

36

Gideon Olarewaju Okunlola, Olusanya Abiodun Olatunji, Richard Olutayo Akinwale, Akash Tariq, Adekunle Ajayi Adelusì.

"Physiological response of the three most cultivated pepper species (*Capsicum* spp.) in Africa to drought stress imposed at three stages of growth and development", *Scientia Horticulturae*, 2017

Publication

<1 %

37

Muhammad Hamzah Saleem, Xiukang Wang, Sajjad Ali, Sadia Zafar et al. "Interactive effects of gibberellic acid and NPK on morpho-physio-biochemical traits and organic acid exudation pattern in coriander (*Coriandrum sativum* L.) grown in soil artificially spiked with boron", *Plant Physiology and Biochemistry*, 2021

Publication

<1 %

38

N Neliyati, L Lizawati, Z Zulkarnain. " The evaluation of sterilization protocol for sprout explants in oil palm (*Jacq.*) tissue culture ", *Journal of Physics: Conference Series*, 2019

Publication

<1 %

39

Nihal M. Elguindy, Galila A. Yacout, Eman F. El Azab, Hala K. Maghraby. "Chemoprotective Effect of *Elettaria Cardamomum* against Chemically induced Hepatocellular Carcinoma in Rats by Inhibiting NF- κ B, Oxidative Stress,

<1 %

and Activity of Ornithine Decarboxylase",
South African Journal of Botany, 2016

Publication

40

Paraserianthes falcataria (L) Nielsen Ekologi
silvikultur dan produktivitas, 2011.

Publication

41

L D Indarwati, E Sulistyyaningsih, B Kurniasih.
"Impact of salicylic acid and biosilica
application on plant growth of shallot under
water deficit", IOP Conference Series: Earth
and Environmental Science, 2021

Publication

42

R Indrasti, J B M Rawung, N R Sudolar, K B
Andri, Sahram, S S Tan. "Farmer institutional
and economic feasibility study on red chilli
pepper farming in Kepung District, Kediri
Regency, East Java Province", IOP Conference
Series: Earth and Environmental Science, 2021

Publication

43

Zulkarnain Z. "Assessment of pollen viability
and germination in Swainsona formosa
(G.Don) J.Thompson", Biospecies, 2019

Publication

44

Vivek Sharma, Richa Salwan, P. N. Sharma,
Arvind Gulati. "Integrated Translatome and
Proteome: Approach for Accurate Portraying
of Widespread Multifunctional Aspects of
Trichoderma", Frontiers in Microbiology, 2017

<1 %

<1 %

<1 %

<1 %

<1 %

45

A Yassi, Kaimuddin, I Ekawati. "Paddy-fish cultivation within an integrated farming system", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

46

Nursanti Nursanti, Novriyanti Novriyanti, Arisa Dwi Indriyani, Nurul Anggraeni, Asrizal Paiman. "Peatlands restorer plant species in the protected forest area", Folia Forestalia Polonica, 2022

Publication

<1 %

47

Nurhaita, N Definiati, U Santoso, S A Akbar, Y L Henuk. " The effect of fermented cocoa pod () husk supplemented with mineral on in vitro digestibility, rumen bacteria population and rumen liquid characteristics ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

48

Saludes, R.B.. "Composting of dairy cattle manure using a thermophilic-mesophilic sequence", Biosystems Engineering, 200710

Publication

<1 %

49

Lina Marija Butkevičienė, Vaida Steponavičienė, Rita Pupalienė, Lina Skinulienė, Rita Čepulienė, Vaclovas Bogužas. "Wheat productivity in different tillage

<1 %

systems after using biostimulants and their mixtures", Zemdirbyste-Agriculture, 2022

Publication

50

Noushina Iqbal, Mehar Fatma, Harsha Gautam, Zebus Sehar, Faisal Rasheed, M.I.R. Khan, Adriano Sofo, Nafees Khan. "Salicylic Acid Increases Photosynthesis of Drought Grown Mustard Plants Effectively with Sufficient-N via Regulation of Ethylene, ABA and Nitrogen-Use Efficiency", Research Square Platform LLC, 2021

Publication

<1 %

51

Parlin H. Sinaga, Marsid Jahari, Usman, Ida Nur Istina, Nana Sutrisna. "Minimum fertilizer for maize cultivation in suboptimal agroecosystem", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

52

R ROSA. "EFFECTS OF TRICHODERMA HARZIANUM AND BORON ON SPRING BROCCOLI", Applied Ecology and Environmental Research, 2019

Publication

<1 %

53

Keiko Yamaji. "The Antifungal Compound Totarol of *Thujopsis dolabrata* var. *hondai* Seeds Selects for Fungi on Seedling Root Surfaces", Journal of Chemical Ecology, 12/05/2007

Publication

<1 %

54 L. E. Francois. "Narrow row cotton (Gossypium hirsutum L.) under saline conditions", Irrigation Science, 1982
Publication <1 %

55 Md. Nasir Hossain Sani, Md. Nazrul Islam, Jasim Uddain, Md. Shah Newaz Chowdhury, Sreeramanan Subramaniam. "Synergistic effect of microbial and nonmicrobial biostimulants on growth, yield, and nutritional quality of organic tomato", Crop Science, 2020
Publication <1 %

56 Michael Danon, Ingrid H. Franke-Whittle, Heribert Insam, Yona Chen, Yitzhak Hadar. "Molecular analysis of bacterial community succession during prolonged compost curing", FEMS Microbiology Ecology, 2008
Publication <1 %

57 Rishi Ram Burlakoti, John Zandstra, Karla Jackson. "Evaluation of epidemics and weather-based fungicide application programmes in controlling anthracnose fruit rot of day-neutral strawberry in outdoor field and protected cultivation systems", Canadian Journal of Plant Pathology, 2014
Publication <1 %

58 "Plant Stress Mitigators", Springer Science and Business Media LLC, 2022
Publication <1 %

-
- 59 "Protective Chemical Agents in the Amelioration of Plant Abiotic Stress", Wiley, 2020
Publication <1 %
-
- 60 Debasis Mitra, Alondra M. Díaz Rodríguez, Fannie I. Parra Cota, Bahman Khoshru et al. "Amelioration of thermal stress in crops by plant growth-promoting rhizobacteria", Physiological and Molecular Plant Pathology, 2021
Publication <1 %
-
- 61 Rafiuddin, A Mollah, E Pratiwi. "Application of various types of local microorganisms to the growth and production of two rice varieties", IOP Conference Series: Earth and Environmental Science, 2020
Publication <1 %
-
- 62 Sustainable Development and Biodiversity, 2014.
Publication <1 %
-
- 63 Zumi Saidah, Harianto, Sri Hartoyo, Ratna Winandi Asmarantaka. "Change on Production and Income of Red Chili Farmers", IOP Conference Series: Earth and Environmental Science, 2020
Publication <1 %
-

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off