

# Junedi\_2022\_IOP\_Conf.\_Ser.\_E arth\_Environ.\_Sci.\_1005\_012013 .pdf

*by Heri Junedi 1*

---

**Submission date:** 09-May-2023 10:54AM (UTC+0700)

**Submission ID:** 2088233716

**File name:** Junedi\_2022\_IOP\_Conf.\_Ser.\_Earth\_Environ.\_Sci.\_1005\_012013.pdf (420.27K)

**Word count:** 3464

**Character count:** 16680

PAPER • OPEN ACCESS


## Effect of coffee husk compost to increase peanut yield in inceptisol

To cite this article: H Junedi *et al* 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1005** 012013

View the [article online](#) for updates and enhancements.

### You may also like

- [Utilization of cacao shell biochar and compost to improve cayenne pepper \(\*Capsicum frutescens\* L.\) in acid upland](#)  
R Ariani, N L Nurida and A Dariah
- [The Long Composting Period Effect of Leaf Shallots on the Compost Quality](#)  
Muhammad Juwanda, Sakhidin, Saparso et al.
- [Carbon storage potential in degraded forests of Kalimantan, Indonesia](#)  
Antônio Ferraz, Sassan Saatchi, Liang Xu et al.




The Electrochemical Society  
Advancing solid state & electrochemical science & technology

243<sup>rd</sup> Meeting with SOFC-XVIII

Boston, MA • May 28 – June 2, 2023

Accelerate scientific discovery!

Learn More & Register



## Effect of coffee husk compost to increase peanut yield in inceptisol

H. Junedi<sup>1</sup>, S Suryanto<sup>1</sup> and H Syukria<sup>2</sup>

<sup>1</sup> Faculty of Agriculture University of Jambi, Jambi, Indonesia

<sup>2</sup> Alumni of Faculty of Agriculture, University of Jambi, Jambi, Indonesia

E-mail: [heri\\_junedi@unja.ac.id](mailto:heri_junedi@unja.ac.id)

**Abstract.** The use of compost as an ameliorant has been shown to improve soil quality and increase crop yields. One of the organic materials that can be used as compost is coffee husk. Besides being able to increase the quality and yield of the plant, the rest of the coffee husk will also pollute the environment if it is not utilized. The aims of this study were to evaluate the effect of coffee husk compost application on Inceptisol and peanut yield. The field experiment was done on an Inceptisol, from March to December 2020. The experiment used a randomized block design with 6 treatment of coffee husk compost with various doses and 4 replications, namely without compost, 10 Mg ha<sup>-1</sup>, 15 Mg ha<sup>-1</sup>, 20 Mg ha<sup>-1</sup>, 25 Mg ha<sup>-1</sup>, and 30 Mg ha<sup>-1</sup>. The results showed that the application of 20 Mg ha<sup>-1</sup> coffee husk compost was the best treatment in optimizing decreasing soil compaction, bulk density, and increasing porosity, while 15 Mg ha<sup>-1</sup> coffee husk compost was the best treatment in increasing peanut yield.

### 1. Introduction

The need for peanuts continues to increase and is not followed by production so that peanut imports are needed in Indonesia. The number of peanut imports in Indonesia increased from 2014 to 2017 and there was a slight decrease in 2018. The number of peanut imports during 2014-2018 was 45,226 Mg, 66,415 Mg, 129,562 Mg, 254,039 Mg, and 244,066 Mg [1].

One of the reasons for the increase in peanut imports is the limited productive agricultural land for producing peanuts due to the conversion of agricultural land for food crops to oil palm plantations and non-agricultural land. One of the soils that has quite a lot of potential is Inceptisol, which is around 20.75 million ha or 37.5% of the total land area of Indonesia [2]. However, Inceptisols generally have low fertility, high sand content, low organic matter and high bulk density. The high soil bulk density describes the compacted soil that is not suitable for peanut plants.

Compacted soil will inhibit root movement in the soil. This will affect the ability of the soil to absorb water, air, and nutrients so that little is available to plants [3]. Soil compaction can be reduced by increasing the organic matter content of the soil.

One source of organic matter is coffee skin, which is a by-product of processed coffee that cannot be utilized optimally. Utilization of coffee husk waste in addition to improving soil properties [4] and increasing crop yields will also help protect the environment [5]-[6]. However, the problem is that coffee husk waste has a very high C/N ratio, 31 [7] so it will take a long time to decompose. Therefore, it is necessary to compost the coffee husk so that it can immediately be used to improve the soil and increase crop yields.



Several studies have shown that coffee husk compost can improve soil properties and increase crop yields. The application of coffee husk compost could increase the production of shallots (*Allium ascalonicum*) [8], coffee [9], red pepper [10], lettuce [11], fruit [12], and chili pepper [13]. The addition of coffee husk compost can also improve soil chemical properties (increase C-organic, total N content, phosphorus and potassium) and soil physical properties (increase aggregate stability and decrease soil density) [14].

Although there have been studies using coffee husk compost in agriculture, no researcher has used peanut plants grown in the Inceptisol. Therefore, the aims of this research are 1) to study whether the application of coffee husk compost can improve soil compaction and at the same time increase peanut crop yields and 2) to find the most appropriate dose to improve soil compaction and produce peanut yield.

## 2. Methods

### 2.1. Compost

The compost is made using a mixture of ingredients: 240 kg of coffee husk, 80 kg of mature cow dung, 4 liters of EM4, 1 kg of shrimp paste and a little brown sugar. Materials are arranged in layers with a height of each layer of  $\pm 15$  cm in a tub measuring 1 m long, 1 m wide, and 1 m high. Each layer consists of a mixture of coffee skin and cow dung in a ratio of 3:1, shrimp paste and brown sugar, then is sprayed with EM4 solution until evenly distributed and then sprayed with water until the mixture is moist. The material that has been arranged in layers is covered, opened every week and stirring and watered. Compost is ready to be applied to the field after 4 months characterized by a blackish color and crumb structure. Husk coffee compost consists of organic-C 30.1%, N-total 1.71%, C/N ratio 17.6%, pH 7.36, and water content 23%.

36

### 2.2. Field experimental design

The field experiment was done on an Inceptisol, from March to December 2020. The soil is low of organic C (2.15%), porosity (45.67%), high bulk density ( $1.31 \text{ g cm}^{-3}$ ) and soil strength ( $18.83 \text{ N cm}^{-2}$  on 310 cm soil layer depth and  $24.92 \text{ N cm}^{-2}$  on a 20 cm soil layer depth).

The treatments were arranged in a randomized block design consisting of six treatments with the addition of coffee husk compost with 6 levels (without compost,  $10 \text{ Mg ha}^{-1}$ ,  $15 \text{ Mg ha}^{-1}$ ,  $20 \text{ Mg ha}^{-1}$ ,  $25 \text{ Mg ha}^{-1}$ , and  $30 \text{ Mg ha}^{-1}$ ) and repeated four times. The compost was applied based on dry weight, spread over the land according to the treatment, mixed and then incubated for two weeks. Peanut seeds were planted with a spacing of 40 cm x 20 cm in plots measuring 3 m x 2 m.

### 2.3. Measurement of soil compaction and plant height

19

Measurements of soil compaction and plant height were carried out at the eighth week. Soil compaction at a depth of 0-10 cm and 10-20 cm was measured using a penetrometer while plant height was measured from the base of the stem to the growing point of the plant.

### 2.4. Sampling and soil analysis

Soil samples were collected from each plot three day prior to harvesting. There were two kinds of soil samples disturbed and undisturbed. Disturbed soil sample to measure soil organic C which was determined by loss of ignition method. An undisturbed soil sample was taken with metal cores of 7.3 cm height and 4.0 cm diameter at the 0-20 cm depth to determine bulk density and porosity.

### 2.5. Sampling and plant analysis

The peanut was harvested 98 days after planting. The pod was cleaned and dry in the sun. The data on yield was recorded from all each plot.

## 2.6. Statistical analysis

The data from the results of research on soil strength and plant height at eight week, bulk density, porosity, organic-C, and yield were analyzed statistically by means of a confidence level ( $\alpha = 5\%$ ). To find out the best compost dose, Duncan's Multiple Spacing test was used at a confidence level ( $\alpha = 5\%$ ).

## 3. Results and discussion

### 3.1. The effect of coffee husk compost on soil organic matter content, bulk density, and porosity

The addition of coffee husk compost into Inceptisol had a significant effect on soil organic carbon content, soil bulk density, and porosity. Further tests using DMRT showed that the higher the dose of compost applied to the soil, the higher the C-organic content and porosity and the lower the soil bulk density but there was a decrease at doses of 25 and 30 Mg ha<sup>-1</sup> of compost. The effect of the application of coffee husk compost on soil organic carbon content, soil bulk density, and porosity are shown in table 1.

**Table 1.** Organic-C content, bulk density and porosity due to the application of coffee husk compost.

Treatment	C-organik (%)	Bulk Density (g cm <sup>-3</sup> )	Porosity (%)
without compost	1.74 a	1.35 b	43.80 a
10 Mg ha <sup>-1</sup> of compost	2.61 c	1.19 ab	46.31 a
15 Mg ha <sup>-1</sup> of compost	2.76 cd	1.13 a	51.68 ab
20 Mg ha <sup>-1</sup> of compost	3.19 d	1.08 a	58.68 c
25 Mg ha <sup>-1</sup> of compost	2.90 cd	1.08 a	54.31 bc
30 Mg ha <sup>-1</sup> of compost	2.18 b	1.23 ab	48.65 ab

Value with the same letters followed in the same column are not significantly different at the 5% level

Table 1 shows that the organic carbon in the treatment without compost was significantly different compared to the coffee husk compost in each treatment. The highest organic matter content was found in the application of coffee husk compost 20 Mg ha<sup>-1</sup> (3.19%) which was significantly different from the treatment with coffee husk compost 10 Mg ha<sup>-1</sup>, 30 Mg ha<sup>-1</sup> and without coffee husk compost.

The lowest bulk density was found in the treatment of coffee husk compost 20 Mg ha<sup>-1</sup>, which was significantly different from without coffee husk compost treatment but not significantly different from the treatment with coffee husk compost 10 Mg ha<sup>-1</sup>, 15 Mg ha<sup>-1</sup>, 20 Mg ha<sup>-1</sup>, 25 Mg ha<sup>-1</sup> and 30 Mg ha<sup>-1</sup>. This is because coffee husk compost has been able to increase soil organic-C. The decrease in bulk density was followed by an increase in soil organic-C, this was in line with research [15] which stated that the application of compost and manure 10 Mg ha<sup>-1</sup> to cassava was able to reduce soil bulk density by 9.82% in Alfisol.

The application of 20 Mg ha<sup>-1</sup> coffee husk compost gave the highest value in soil porosity. The high value of porosity is due to the high content of soil organic-C. According to [16] the addition of organic material in the form of compost will increase the porosity of the soil so that the total value of the pore space is high.

Adding organic matter to the soil will increase the organic-C content. The increase in organic-C affects the value of bulk density and porosity. The less organic-C contained in the soil, the higher the bulk density and the lower the porosity. The results of this study are the same as those of [17] which stated that increasing soil organic matter content will reduce soil bulk density and increase porosity.

### 3.2. <sup>15</sup> effect of coffee husk compost on soil compaction

The effect of the application of coffee husk compost on soil compaction in the eighth week at a depth of 0-10 cm and a depth of 10-20 cm are shown in table 2.

<sup>19</sup> **Table 2.** Soil compaction at a depth of 0-10 cm and 10-20 cm at week 8 due to the application of coffee husk compost.

Treatment	Soil compaction (Nm <sup>-2</sup> )			
	Depth 0-10 cm		Depth 10-20 cm	
Without compost	9.67	a	21.17	a
10 Mg ha <sup>-1</sup> of compost	8.25	b	17.08	bc
15 Mg ha <sup>-1</sup> of compost	6.33	c	15.75	cd
20 Mg ha <sup>-1</sup> of compost	4.67	d	11.17	e
25 Mg ha <sup>-1</sup> of compost	5.92	cd	12.50	de
30 Mg ha <sup>-1</sup> of compost	9.25	4 b	20.08	ab

Value with the same letters followed in the same column are not significantly different at the 5% level

Based on Table 2, it can be seen that overall soil compaction at a depth of 10 cm and 20 cm has the same tendency, which occurs at the lowest rate of 20 ton ha<sup>-1</sup> of coffee husk compost, which is significantly different from all treatments except for the treatment of coffee husk compost 25 tons ha<sup>-1</sup>. According to [18] compost is able to improve several physical properties of the soil, namely increasing the ability of the soil to hold water, reducing soil compaction and increasing soil porosity so that it can help increase plant growth with good root growth.

The level of soil compaction can be characterized by high bulk density and low porosity<sup>22</sup>. Soil compaction will continue to increase with increasing soil depth. It occurred due to the lower levels of organic-C in the soil. In addition, the dose of coffee husk compost up to a depth of 20 cm also causes high soil compaction. The increase in soil compaction is also due to the decreasing organic-C content so that the bulk density content increases with increasing soil depth.

### <sup>3</sup> 3. The effect of coffee husk compost on height and yield of peanuts

<sup>42</sup> results of variance showed that the application of coffee husk compost on Inceptisol had a very significant effect on plant height and peanut yield. The effect of coffee husk compost on height and yield of peanuts are shown in table 3.

<sup>33</sup> **Table 3.** The effect of coffee husk compost on height and yield of peanuts.

Treatment	Height of peanuts (cm)		Yield of peanuts (Mg ha <sup>-1</sup> )	
Without compost	62.13	a	1.69	a
10 Mg ha <sup>-1</sup> of compost	66.78	b	1.89	ab
15 Mg ha <sup>-1</sup> of compost	68.69	bc	1.97	b
20 Mg ha <sup>-1</sup> of compost	72.25	d	2.03	b
25 Mg ha <sup>-1</sup> of compost	70.00	cd	1.98	b
30 Mg ha <sup>-1</sup> of compost	64.19	4 a	1.78	ab

Value with the same letters followed in the same column are not significantly different at the 5% level

Table 3 shows that the 20 ton ha<sup>-1</sup> treatment was the best treatment in increasing plant height. The treatment of 20 ton ha<sup>-1</sup> coffee husk compost showed a significant difference to all treatments. This is in line with the research of [19] that coffee husk compost was able to restore soil fertility and increase the growth of coffee seedling plant height compared to controls.



The treatment of the coffee husk compost 15 to 29 ha<sup>-1</sup> was significantly different from the treatment without compost, the coffee husk compost 10 tons ha<sup>-1</sup>, and the coffee husk compost 30 tons ha<sup>-1</sup> but not significantly different from the coffee husk compost 20 tons ha<sup>-1</sup> and the coffee husk compost 25 tons ha<sup>-1</sup>. This result shows that the application of 15 ton ha<sup>-1</sup> of coffee husk compost has been able to increase the yield of peanuts. This is presumably because the amount of organic matter in the 15 ton ha<sup>-1</sup> treatment was able to increase the yield of peanuts.

The ability of the soil to absorb nutrients optimally increases due to the application of the coffee husk compost. [20] stated that compost can increase nutrients in the soil. This is in line with [7] stated that husk coffee compost can increase Phosphorus and Potassium. Composting can also create suitable soil conditions for plant growth by increasing soil organic matter, reducing soil compaction. These conditions support plant growth to have optimal growth and productivity.

#### 4. Conclusions

Based on the results of the study it can be concluded as follows:

1. The application of coffee husk compost affects the organic C content, bulk density, porosity, soil compaction, height and yield of peanuts
2. The best treatment in reducing soil compaction from 9.67 N cm<sup>-1</sup> to 4.67 N cm<sup>-2</sup> is coffee husk compost 20 Mg ha<sup>-1</sup>
3. The best treatment in increasing height and yield of peanuts is husk coffee compost 15 Mg ha<sup>-1</sup>

#### References

- [1] Badan Pusat Statistik 2019 Statistik perkembangan produksi, konsumsi, dan impor kacang tanah di Indonesia. Diunduh dari <http://www.bps.go.id>
- [2] Muyassir, Sufardi, dan I Saputra 2012 Perubahan sifat fisika inceptisol akibat perbedaan jenis dan dosis pupuk organik *Jurnal Lentera* **12**(1)
- [3] Arthur E, Cornelis W, and Razzaghi F Compost amendment to sandy soil affects soil properties and greenhouse tomato productivity *Compost Science & Utilization* **20** 215
- [4] Insam H and de Bertoldi M 2007 Microbiology of composting process *Compost Science and Technology*, ed Golke C, Bidlingmaier, de Bertoldi M, and Diaz L (Eds.) pp 25
- [5] Murthy PS and Maidu M 2012 Sustainable management of coffee industry by-product and value addition-a review of Resour. Conserv. Recycle. **66** 45
- [6] Zoca SM, Penn CJ, Rosolem CA, Alves AR, Neto LO, and Martins MM 2014 Coffee processing residues as a soil potassium amendment *Int.J Recycl Org Waste Agric* **3** 155
- [7] Putra RE, Hutami R, Suantika G, and Rosmiati M 2017 Application of compost produced by bioconversion of coffee husk by black soldier fly larvae (*Hermetia illucens*) as solid fertilizer to lettuce (*Lactuca sativa* var. *crispa*) impact to harvested biomass and utilization of Nitrogen, Phosphor, and Potassium *Green Tech* **8**(1) 20
- [8] Simanjuntak, Ratna R and Purba E 2013 Respon pertumbuhan dan produksi bawang merah (*allium ascalonicum* L.) terhadap pemberian pupuk NPK dan kompos kulit buah kopi *Jurnal Online Agroekoteknologi* **3**(1) 362
- [9] Dzong NA, Dzong TT, Khanh VTP 2012 Evaluation of Coffee Husk Compost for Improving Soil Fertility and Sustainable Coffee Production in Rural Central Highland of Vietnam *Resources and Environment* **3** (4) 77
- [10] Berlian Z, Syarifah S and Sari DS 2015 Pengaruh pemberian limbah kulit kopi (*Coffea robusta* L.) terhadap pertumbuhan cabai (*Capsicum annum* L.) *J Biota* **1** (1) 22
- [11] Getachew G and Mureta D 2017 Optimization of compost maturity of coffee waste mixed with agricultural wastes and evaluation of their effect on growth of lettuce (*Lactuca sativa*) *Journal of Natural Science Research* **7** (8) 82
- [12] Fatoni A, Muhfahroyin, Sujarwanta A, Sutanto A, Widowati H 2020 The effectiveness of coffee leather organic fertilizer on the productivity of red spinach (*Amaranthus tricolor* L.) *Jurnal Penelitian Pertanian Terapan* **20** (3) 198

- [13] Adnan, Basri AB, Azis A, Idawanni and Iswoyo H 2021 Application of coffee husk compost and EM4 on growth and yield of chili pepper (*Capsicum frutescens* L.) IOP Conf. Ser.: Earth Environ. Sci. **807** 042040
- [14] Valentia FV, Listyarini E, Prijono S 2015 Aplikasi kompos kulit kopi untuk perbaikan sifat kimia dan fisika tanah inceptisol serta meningkatkan produksi brokoli *Jurnal Tanah dan Sumberdaya Lahan* **2(1)** 147
- [15] Prasetyo A, Utomo WH dan Listyorini E 2014 Hubungannya sifat fisik tanah, perakaran dan hasil ubi kayu tahun kedua pada alfisol jatikerto akibat pemberian pupuk organik dan anorganik (NPK) *Jurnal Tanah dan Sumberdaya Lahan* **1(1)** 27
- [16] Hasibuan ASZ 2015 Pemanfaatan bahan organik dalam perbaikan beberapa sifat tanah pasir Pantai Selatan Kulon Progo *Planta Tropika Journal of Agro Science* **3(1)** 31
- [17] Junedi H, Zurhalena, Mahbub IA 2013 Short-term effect of china violet compost on soil properties of ultisol and peanut yield *International Journal on Advanced Science Engineering Information Technology* **3 (1)** 50
- [18] Widodo KH dan Kusuma Z 2018 Pengaruh kompos terhadap sifat fisik tanah dan pertumbuhan tanaman jagung di Inceptisol *Jurnal Tanah dan Sumberdaya Lahan* **5(2)** 959
- [19] Falahuddin I, Raharjeng ARP, Harmeni L 2016 Pengaruh pupuk organik limbah kulit kopi (*coffea arabica* L.) terhadap pertumbuhan bibit kopi *Jurnal Bioilmi* **2(2)** 108
- [20] Falahuddin I, Raharjeng ARP, Harmeni L 2016 Pengaruh pupuk organik limbah kulit kopi (*coffea arabica* L.) terhadap pertumbuhan bibit kopi *Jurnal Bioilmi* **2(2)** 108

#### Acknowledgments

The Authors would like to say thank you to all of the people who were involved in this research for their technical help during field work, contribution and support.



ORIGINALITY REPORT

---

20%  
SIMILARITY INDEX

%  
INTERNET SOURCES

20%  
PUBLICATIONS

%  
STUDENT PAPERS

---

PRIMARY SOURCES

---

- 1

St Sabahannur, Suraedah Alimuddin, St Subaedah, Nurul Wahdaniya, Firda Yunita Faharuddin. "The Effect of Cocoa Butter Substitute (CBS) Produced from Palm Kernel Oil and Virgin Coconut Oil (VCO) on The Quality of Milk Chocolate", Research Square Platform LLC, 2023

Publication

1 %
  - 2

Franzluebbers, A.J.. "Surface-soil responses to paraplowing of long-term no-tillage cropland in the Southern Piedmont USA", Soil & Tillage Research, 200710

Publication

1 %
  - 3

Irfan Nofriandi, Faiz Barchia, Fahrurrozi Fahrurrozi, Bilman W. Simanihuruk, Entang Inorah Sukarjo. "Stem Cutting Growth of Red Dragon Fruit (*Hylocereus costaricensis*) due to Aplication of Cow Manure and KCl in Andosol", TERRA : Journal of Land Restoration, 2021

Publication

1 %
-

4

Bambang Irawan, Franz Gruber, Reiner Finkeldey, Oliver Gailing. "Linking indigenous knowledge, plant morphology, and molecular differentiation: the case of ironwood (*Eusideroxylon zwageri* Teijsm. et Binn.)", Genetic Resources and Crop Evolution, 2015

Publication

1 %

5

D. Dadi, G. Daba, A. Beyene, P. Luis, B. Van der Bruggen. "Composting and co-composting of coffee husk and pulp with source-separated municipal solid waste: a breakthrough in valorization of coffee waste", International Journal of Recycling of Organic Waste in Agriculture, 2019

Publication

1 %

6

Irfan Rusdi, Abdul Rauf, Supriadi Supriadi, Benny Hidayat. "Application of biochar from palm oil plants residues on physical properties of Ultisol", AGRITROPICA : Journal of Agricultural Sciences, 2020

Publication

1 %

7

Lukman Lukman. "PENGARUH PENGGUNAAN PUPUK ORGANIK CAIR NIRA AREN [*Arenga pinnata* (WURMB) MERR] TERHADAP PERTUMBUHAN BIBIT TANAMAN CENGKEH (*Syzygium aromaticum*. L)", Jurnal Agrotek Tropika, 2022

Publication

1 %

8

"Medicinal Plants and Fungi: Recent Advances in Research and Development", Springer Science and Business Media LLC, 2017

Publication

1 %

9

D. Purkayastha, S. Sarkar. "Sustainable waste management using black soldier fly larva: a review", International Journal of Environmental Science and Technology, 2021

Publication

1 %

10

Apri Salfiani, Asri Pirade Paserang. "PENGARUH KOMBINASI IAA (Indole-3-Acetic Acid) DAN BAP (6-Benzylaminopurine) TERHADAP INISIASI TANAMAN VANILI (*Vanilla planifolia* Andrews)", Biocelebes, 2022

Publication

1 %

11

Aswari Aswari, Santi R, Lestari T. "PEMANFAATAN LIMBAH KELAPA SAWIT PADA BEBERAPA VARIETAS SORGUM (*Sorghum bicolor* [L.] Moench.) YANG DIBUDIDAYAKAN DI SELA KELAPA SAWIT", Jurnal Ilmiah Hijau Cendekia, 2022

Publication

1 %

12

Edza Wikurendra, Globila Nurika, Novera Herdiani, Yauwan Tobing Lukiyono. "Evaluation of the Commercial Bio-Activator and a Traditional Bio-activator on Compost Using Takakura Method", Journal of Ecological Engineering, 2022

1 %

13

Cristofora Hanny Wijaya, Manik Harda, Bunga Rana. " Diversity and Potency of spp. Grown in Indonesia ", IntechOpen, 2020

Publication

---

14

İlknur Gümüş, Cevdet Şeker. "Effects of Spent Mushroom Compost on Physicochemical Properties of Degraded Soil", Copernicus GmbH, 2016

Publication

---

15

K. W. Waldron. "Composting of food-chain waste for agricultural and horticultural use", Handbook of waste management and co-product recovery in food processing, 2009

Publication

---

16

Mohd Huzairi Mohd Zainudin, Aisyah Zulkarnain, Ain Sahira Azmi, Shalini Muniandy et al. "Enhancement of Agro-Industrial Waste Composting Process via the Microbial Inoculation: A Brief Review", Agronomy, 2022

Publication

---

17

Ouedraogo, E.. "Use of compost to improve soil properties and crop productivity under low input agricultural system in West Africa", Agriculture, Ecosystems and Environment, 200105

Publication

---

1 %

1 %

1 %

1 %

1 %

18

Anisah Mohammed, Mohamad Amir Shah Yusop, Nur Ain Zubaidah Amran. "Chapter 92 Utilization Effects of Asystasia intrusa Compost Toward the Growth Performance of Elaeis guineensis Seedling", Springer Science and Business Media LLC, 2018

Publication

1 %

19

Hiroaki Ikeda, Kiyoshi Okutomi. "Effects of human trampling and multispecies competition on early-phase development of a tread community", Ecological Research, 1990

Publication

<1 %

20

Amalia Michail, Philothei Sigala, Spyros Grigorakis, Dimitris P. Makris. "Kinetics of Ultrasound-Assisted Polyphenol Extraction from Spent Filter Coffee Using Aqueous Glycerol", Chemical Engineering Communications, 2015

Publication

<1 %

21

Bonifasius Nahak. "Pertumbuhan dan Hasil Kacang Hijau (*Vigna radiata* L.) dalam Tumpangsari dengan Jagung (*Zea mays* L.) pada Penanaman Ketiga Di Tanah Entisol Semi Arid yang Mengandung Residu Kompos Biochar dan Pernah Ditanami Jagung Secara Monokultur", Savana Cendana, 2021

Publication

<1 %



22

C.P. Kushwaha, S.K. Tripathi, K.P. Singh. "Soil organic matter and water-stable aggregates under different tillage and residue conditions in a tropical dryland agroecosystem", *Applied Soil Ecology*, 2001

Publication

<1 %

23

Suria Darma Tarigan, Sunarti, Kerstin Wiegand, Claudia Dislich, Bejo Slamet, Johannes Heinonen, Katrin Meyer. "Mitigation options for improving the ecosystem function of water flow regulation in a watershed with rapid expansion of oil palm plantations", *Sustainability of Water Quality and Ecology*, 2016

Publication

<1 %

24

Darso Sugiono, Vera O Subardja, Briljan Sudjana. "Peningkatan Kualitas Fisika Tanah Guna Efisiensi Air Melalui Pengkayaan Media Tanam Dengan Kompos Plus Pada Budidaya Tanaman Jagung Manis", *AGROSAINSTEK: Jurnal Ilmu dan Teknologi Pertanian*, 2018

Publication

<1 %

25

Dessy A Ralahalu, Rhony E Ririhena, Abdul K Kilkoda. "PEMBERIAN PUPUK ORGANIK DAN JARAK TANAM UNTUK PERTUMBUHAN DAN HASIL TANAMAN BAWANG MERAH (*Allium ascalonicum* L.)", *JURNAL BUDIDAYA PERTANIAN*, 2017

<1 %

26

Ryan, J.. "Response to residual and currently applied phosphorus in dryland cereal/legume rotations in three Syrian Mediterranean agroecosystems", European Journal of Agronomy, 200802

Publication

---

27

Lubanga Lunze, Mathew M., Robin Buruchara, Michael A., Nsharwasi Lon, Gideon O., Mulangwa Ngongo, Idupulapati Rao. "Chapter 5 Integrated Soil Fertility Management in Bean-Based Cropping Systems of Eastern, Central and Southern Africa", IntechOpen, 2012

Publication

---

28

T. Hellemans, S. Landschoot, K. Dewitte, F. Van Bockstaele, P. Vermeir, M. Eeckhout, G. Haesaert. "Impact of Crop Husbandry Practices and Environmental Conditions on Wheat Composition and Quality: A Review", Journal of Agricultural and Food Chemistry, 2018

Publication

---

29

M. Sameni, A. Tajabadi Pour. "Effect of root refuse of liquorice on different characteristics of soils from southern regions of Iran. III. Structure and hydraulic conductivity of soils under saline sodic conditions",

<1 %

<1 %

<1 %

<1 %

# Communications in Soil Science and Plant Analysis, 2011

Publication

30

Omar Latifah, Osumanu Haruna Ahmed, Nik Muhamad Abdul Majid. " Short Term Enhancement of Nutrients Availability in L. Cultivation on an Acid Soil Using Compost and Clinoptilolite Zeolite ", Compost Science & Utilization, 2016

Publication

<1 %

31

Rivero, C.. "Influence of compost on soil organic matter quality under tropical conditions", Geoderma, 200412

Publication

<1 %

32

"Fundamental and Applied Sciences in Asia", Springer Science and Business Media LLC, 2022

Publication

<1 %

33

Catur Herison, Rustikawati Rustikawati, Hasanudin Hasanudin, Usman K.J. Suharjo, Merakati Handajaningsih, Nursalim Nursalim. "Manually Feed NPK Fertigation Improves Growth and Yield of Chili Pepper Hybrids in Ultisol", Akta Agrosia, 2020

Publication

<1 %

34

Dedik Budianta, Adipati Napoleon, Akbar Paripurna, E Ermatita. "Growth and production of soybean (Glycine max (L.) Merrill)

<1 %

with different fertilizer strategies in a tidal soil from South Sumatra, Indonesia", Spanish Journal of Soil Science, 2019

Publication

---

35

Hemmatollah Pirdashti, Alaleh Motaghian, Mohammad Ali Bahmanyar. "EFFECTS OF ORGANIC AMENDMENTS APPLICATION ON GRAIN YIELD, LEAF CHLOROPHYLL CONTENT AND SOME MORPHOLOGICAL CHARACTERISTICS IN SOYBEAN CULTIVARS", Journal of Plant Nutrition, 2010

Publication

---

36

Yuxue Zhang, Zuoxin Tang, Yeming You, Xiaowei Guo, Chuanjing Wu, Shirong Liu, Osbert Jianxin Sun. "Differential effects of forest-floor litter and roots on soil organic carbon formation in a temperate oak forest", Soil Biology and Biochemistry, 2023

Publication

---

37

Margarida Arrobas, Peltier Aguiar, M. Ângelo Rodrigues. "A comparison of a pasture ley with a maize monoculture on the soil fertility and nutrient release in the succeeding crop", Archives of Agronomy and Soil Science, 2015

Publication

---

38

C. D. Tsadilas, I. K. Mitsios, E. Golia. "Influence of Biosolids Application on Some Soil Physical

<1 %

<1 %

<1 %

<1 %

# Properties", Communications in Soil Science and Plant Analysis, 2005

Publication

39

Henni Elfandari, Betari Safitri. "Effect of Planting Media Composition on Growth of Chrysanthemum spp.", JURNAL AGROTROPIKA, 2022

Publication

<1 %

40

Johannes Simbolon, Bilman Wilman Simanihuruk, Bambang Gonggo Murcitro, Herry Gusmara, Eko Suprijono. "PENGARUH SUBTITUSI PUPUK N SINTETIK DENGAN LIMBAH LUMPUR SAWIT TERHADAP PERTUMBUHAN DAN HASIL JAGUNG MANIS", Jurnal Ilmu-Ilmu Pertanian Indonesia, 2018

Publication

<1 %

41

Mariangela Diacono. "Long-term effects of organic amendments on soil fertility. A review", Agronomy for Sustainable Development, 2009

Publication

<1 %

42

Nerty Soverda, Evita Evita, Elly Indraswari. "Weeds Pressuring on the Growth and Yield of Soybean Plants By Giving an Acasia Organic Mulch", Current Agriculture Research Journal, 2014

Publication

<1 %



---

Exclude quotes      Off

Exclude matches

< 3 words

Exclude bibliography      Off